

7. Literaturverzeichnis

1. Amico, S.; Liehn, J.C.; Desoize, B.; Larbre, H.; Deltour, G.; Valeyre, J.: Comparison of phosphatase isoenzymes PAP and PSA with bone scan in patients with prostate carcinoma.
Clin Nucl Med 16 (1991), 643-648
2. Aydiner, A.; Topuz, E.; Disci, R.; Yasasever, V.; Dincer, M.; Dincol, K.; Bilge, N.: Serum tumor markers for detection of bone metastasis in breast cancer patients.
Acta Oncol 33 (1994), 181-186
3. Berndt, L.A.; Sieber, E.; Berndt, R.: Morbidität und Mortalität bösartiger Neubildungen der Brustdrüse.
Zeitschr Ärztliche Fortbildung 73 (1979), 1123-1129
4. Berruti, A.; Torta, M.; Piovesan, A.; Raucci, C.A.; Orlandi, F.; Panero, A.: Biochemical picture of bone metabolism in breast cancer patients with bone metastases.
Anticancer Res 15 (1995), 2871-2876
5. Berruti, A.; Osella, G.; Raucci, C.A.; Roncari, A.; Dogliotti; L.: Transient increase in total serum alkaline phosphatase predicts radiological response to systemic therapy in breast cancer patients with osteolytic and mixed bone metastases.
Oncology 50 (1993), 218-221
6. Blijlevens, N.M.; Oosterhuis, W.P.; Oosten, H.R.; Mulder, N.H.: Clinical value of TPS, CEA and CA 15-3 in breast cancer patients.
Anticancer Res 15 (1995), 2711-2716
7. Body, J.J.; Dumon, J.C.; Gineyts, E.; Delmas, P.D.: Comparative evaluation of markers of bone resorption in patients with breast cancer-induced osteolysis before and after bisphosphonate therapy.
Br J Cancer 75 (1997), 408-412

8. Bohmert, H.; Eiermann, W.; von Hesler, F.W.; Drzeska, W.; Willich, N.; Lindner, H.:
Interdisziplinäres Vorgehen in der Primärtherapie des Mammakarzinoms.
In: Schriftenreihe des Tumorzentrum München mit Empfehlungen zur Diagnostik,
Therapie und Nachsorge onkologischer Erkrankungen; Tumorzentrum München
(1991), 11-28

9. Broyn, J.; Froyen, T.:
Evaluation of routine follow-up after surgery for breast carcinoma.
Acta Chem Scand 148 (1982), 401-404

10. Brule, G.:
Co-operative clinical study of 178 patients treated with "Novaldex".
In: The hormonal control of breast cancer.
Hrsg: Brule, G.
ICI Pharmaceutical Division Ltd Alderly Park, Macclesfield, Cheshire (1978),
35-39

11. Burlina, A.; Rubin, D.; Secchiero, S.; Sciacovelli, L.; Zaninotto, M., Plebani, M.:
Monitoring skeletal cancer metastases with the bone isoenzyme of tissue
unspecific alkaline phosphatase.
Clin Chem Acta 226 (1994), 151-158

12. Caffier, H.; Brandau, H.:
Serum tumor markers in metastatic breast cancer and course of disease.
Cancer Detect Prev 6 (1983), 451

13. Carey, P.O.; Lippert, M.C.:
Treatment of painful prostatic bone metastases with oral etidronate disodium.
J Urol 32 (1988), 403-407

14. Coleman, R.E.; Mashiter, G.; Whitaker, K.B.; Moss, D.W.; Rubens, R.D.; Fogelman, I.:
Bone scan flare predicts successful systemic therapy for bone metastases.
J Nucl Med 29 (1988), 1354-1389
15. Coleman, R.E.; Rubens, R.D.:
Bone metastases and breast cancer.
Cancer Treat Rev 12 (1985), 251-270
16. Coleman, R.E.; Whitaker, K.B.; Moss, D.W.; Mashiter, G., Fogelman, I.; Rubens, R.D.:
Biochemical prediction of response of bone metastases to treatment.
Br J Cancer 58 (1988), 205-210
17. Cooper, E.H.; Forbes, M.A.; Hancock, A.K.; Parker, D.; Laurence, V.:
Serum bone alkaline phosphatase and CA 549 in breast cancer with bone metastases.
Biomed Pharmacother, 46 (1992), 31-36
18. Cooper, E.H.; Purves, D.A.; Yerna, M.J.:
Tandem Ostase - An IRMA for bone alkaline phosphatase evaluated in breast and prostate cancer.
In: Current tumor diagnosis: applications, clinical relevance, research, trends, cancer of the lung – state and trends in diagnosis and therapy.
Hrsg: Klapdor, R.
Zuckerschwertd München, Bern, Wien, New York, (1994)
19. Creutzig, H.:
Bedeutung der Knochenszintigraphie für die Nachsorge des Mammakarzinoms.
Deutsche Med Wochenschrift 111 (1986), 1860-1861

20. Crippa, F.; Bombardieri, E.; Seregni, E.; Castellani, M.R.; Gasparini, M.; Maffioli, L.; Pizzichetta, M.; Buraggi, G.L.:
Single determination of CA 15-3 and bone scintigraphy in the diagnosis of skeletal metastases of breast cancer.
J Nucl Med, 36 (1992), 52-55
21. Crivellari, D.; Price, K.N.; Hagen, M.; Goldhirsch, A.; Gelber, R.D.; Castiglione, M.; Coates, A. S.; Rudenstam, C.M.; Collins, J.; Lindtner, J.:
Routine tests during follow-up of patients after primary treatment for operable breast cancer.
Ann Oncol, 6 (1995), 769-776
22. Crombach, G.:
Überlegungen zum rationellen Einsatz von Tumormarkern beim Mammakarzinom.
Schweiz Rundsch Med Prax, 87 (1998), 589-594
23. Demers L.M.; Costa, L.; Chinchilli, V.M.; Gaydos, L.; Curley, E.; Lipton, A.:
Biochemical markers of bone turnover in patients with metastatic bone disease.
Clin Chem 41 (1995), 1489-1494
24. De Waard, F.:
Epidemiology of breast cancer – a review.
Cancer Clin Oncol 19 (1983), 1671-1676
25. Davies, J.; Trask, C.; Souhami, R.L.:
Effect of mithramycin on widespread painful bone metastases in cancer of the breast.
Cancer Treat Rep 63 (1979), 1835-1838
26. Desoize, B.; Veiler, V.; Pourny, C.; Comoe, L.; Jardillier, J.C.:
Isoenzymes of alkaline and acid phosphatases as bone metastasis marker in breast cancer patients.
Anticancer Res 9 (1989), 1105-1109

27. Desoize, B.; Pourny, C.; Amico, S.; Larbre, H.; Jardillier, J.C.:
Evaluation de deux isoenzymes seriques des phosphatases comme marqueurs
des metastases osseuses.
Bull Cancer (Paris) 77 (1991), 1211-1221
28. Diel,I.J.; Costa, S.D.; Kaufmann, M.; Schmid, H.; Bastert, G.:
Systemische Therapie des ossär metastasierten Mammakarzinoms (OMM).
In: Knochenmetastasen: Pathophysiologie, Diagnostik und Therapie.
Hrsg.: Wüster,C.; Ziegler, R.
Zuckerschwerdt, München, Bern, Berlin
29. Ebeling, K.; Nischan, P.:
Mammakarzinom-Epidemiologie und Prävention.
Zeitschr Ärztliche Fortbildung 85 (1991), 956
30. Engel, K.; Schmid, H.; Hanke, J.; Kaufmann, M.; Müller, A.:
CA 15-3 und CEA als Tumormarker zur Rezidivdiagnostik beim Mammakarzinom.
Geburtsh Frauenheilk 48 (1988), 309-312
31. Fleisch,H.:
Bisphosphonate bei Knochenerkrankungen.
H. Huber Verlag, Bern (1997)
32. Francini, G.; Montagnani, M.; Petrioli, R.; Paffetti, P.; Marsili, S.; Leone, V.:
Comparison between CEA, TPA, CA 15/3 and hydroxyproline, alkaline
phosphatase, whole body retention of 99mTc MDP in the follow-up of bone
metastases in breast cancer.
Int J Biol Markers 5 (1990), 65-72
33. Galasko, C.S.B.:
Skeletal metastases and mammary cancer.
Ann R Coll Surg Engl 50 (1972), 3-28

34. Galasko, C.S.B.:
The pathological basis for skeletal scintigraphy.
J Bone Joint Surg Br 57-B (1975), 353-359
35. Gang, Y.; Adachi, T.; Ohkura, H.; Yamamoto, H.; Mizuguchi, Y.; Abe, K.:
CA 15-3 is present as a novel tumor marker in the sera of patients with breast cancer and other malignancies.
Gan To Kagaku Ryoho (Jap J Cancer Chemother) 12 (1985), 2379-2386
36. Gerber, B.; Wilken, H.:
Nachsorge beim Mammakarzinom.
Ärztebl Mecklenburg-Vorpommern 12 (1992), 1250-1258
37. Go, V.L.:
Carcinoembryonales antigen: Clinical application.
Cancer 37 (1976), 562-566
38. Goldenberg, D.M.; Neville, M.; Carter, A.C.:
CEA (carcinoembryonic antigen): its role as a marker in the management of cancer.
J Cancer Clin Oncol 101 (1981), 239-242
39. Gomez, B.J.; Ardakani, S.; Ju, J.; Jenkins, D.; Cerelli, M.J.; Daniloff, G.Y.; Kung, V.T.:
Monoclonal antibody assay for measuring bone-specific alkaline phosphatase activity in serum.
Clin Chem 41 (1995), 1560-1566
40. Harris, J.; Hellmann, S.:
Observations on survival curve analysis with particular reference to breast cancer treatment.
Cancer 57 (1986), 925-928

41. Harris, H.:
The human alkaline phosphatases: what we know and what we don't know.
Clin Chim Acta 186 (1989), 133

42. Hasert, V.:
Diagnostische Verfahren zur Ergänzung der Mammographie.
Z ärztl Fortbild 85 (1991), 1001-1006

43. Hayes, F.; Zurawsky, V.R.; Kufe, D.W.:
Comparison of circulating CA 15-3 and carcinoembryonic antigen levels in patients with breast cancer.
J Clin Oncol 4 (1986), 1542-1550

44. Hellriegel, K.P.; Schulz, K.D.:
Nachsorge bei Mammakarzinom-Patientinnen. Empfehlungen einer Konsensus-Tagung, Berlin 23. - 24.02. 1995.
Oncology 1 (1995), 405-412

45. Höffken, K.; Kempf, H.; Miller, A.A.; Miller, B.; Schmidt, C.G.; Faber, P.; Kley, H.K.:
Aminoglutethimide without hydrocortisone in the treatment of postmenopausal patients with advanced breast cancer.
Cancer Treat Rep 70 (1986), 1153-1157

46. Hossfeld, K.; Kleeberg, U.R.:
Gedanken zur Nachsorge bei Patientinnen mit Mammakarzinom.
Deutsches Ärztebl 90 (1993), 437-439

47. Hözel, D.; Thieme, C.H.:
Die Skelettszintigraphie in der Nachsorge des Mammakarzinoms. Statistische und epidemiologische Gesichtspunkte.
Deutsche Med Wsch 111 (1986), 1191-1199

48. Hortobagyi, G.N.; Libshitz, H.I.; Seabold, J.E.:
Osseous metastases of breast cancer. Clinical, biochemical, radiographic and scintigraphic evaluation of response to therapy.
Cancer 53 (1984), 577-582
49. Horton, J.:
Follow-up of breast cancer patients.
Cancer 53 (1984), 790-797
50. Kamby, C.; Vejborg, I.; Daugard, S.; Guldhammer, B.; Dirksen, H.; Rossing, N.:
Clinical and radiologic characteristics of bone metastases in breast cancer.
Cancer 60 (1987), 2524-2531
51. Kamby, C.; Egsmose, C.; Søletormos, G.; Dombernowsky, P.:
The diagnostic and prognostic value of serum bone Gla protein (osteocalcin) in patients with recurrent breast cancer.
Scand J Clin Lab Invest 53 (1993), 439-446
52. Khandekar, J. D.:
Recommendations on follow-up of breast cancer patients following primary therapy.
Sem Surg Oncol 12 (1996), 346-351
53. Kleist, v. S.:
Die Rezidiverkennung von Mammakarzinomen mit Hilfe von Tumormarkerbestimmungen.
In: Aktuelle Aspekte in der Diagnostik des Mammakarzinoms.
Hrsg: Jonat, W.
Bücherei des Frauenarztes 31 (1989), 57

54. Kloskowski, S.; Ebeling, K.:
Nutzen der monatlichen Selbstuntersuchung der Brust für die Früherkennung des Mammakarzinoms.
Arch Geschwulstforsch 60 (1990), 377-382
55. Kreienberg, R.:
Allgemeine und spezifische Laborparameter im Rahmen der Tumornachsorge bei gynäkologischen Malignomen und bei Mammakarzinomen.
Gynäkologe 22 (1989), 55-62
56. Lamerz, R.; Fateh-Moghadam, A.:
Carcinofetale Antigen. II. Carcinoembryonales Antigen (CEA).
Klin Wochenschr 53 (1975), 193-203
57. Langlands, A.; Pocock, S.; Kerr, G.; Gore, S.M.:
Long-term survival of patients with breast cancer: a study of the curability of the disease.
Br Med J 2 (1979), 1247-1251
58. Leonhardt, A.:
Konzeption der programmierten und standardisierten Mammakarzinom-Nachsorge.
Münch Med Wochenschr 118 (1976), 297-302
59. Lerner, H.J.; Band, P.R.; Israel, L.; Leung, B.S.:
Phase II study of tamoxifen: report of 74 patients with stage IV breast cancer.
Cancer Treat Rep 60 (1976), 1431-1435
60. Liberati, A.:
The GIVIO trial on the impact of follow-up care on survival and quality of life in breast cancer patients.
Ann Oncol 6 (1995), 41-46,

61. Loprinzi, C.L.; Ahmann, D.L.:
Carcinoembryonic antigen. A routine test in patients with breast carcinoma ?
Arch Int Med 146 (1986), 2125-2126
62. Lorente, J.A.; Morote, J.:
Behavior of bone alkaline phosphatase (BAP) determined with
immunoradiometric assay in metastatic prostate cancer.
Int J Biol Markers 9 (1994), 145
63. Mackintosh, J.; Simes, J.; Raghavan, D.; Pearson, B.:
Prostatic cancer with bone metastases: serum alkaline phosphatase (SAP) as a
predictor of response and the significance of the SAP "flare".
Br J Urol 66 (1990), 88-93
64. Marchai, P.; Santini, D.; Bianco, V.; Chiodini, S.; Reale, M. G.; Simeoni, F.;
Marchei, G. G.; Vecchione, A.:
Serum ostase in the follow-up of breast cancer patients.
Anticancer Res 15 (1995), 2217-2222
65. Martinetti, A.; Bajetta, E.; Seregni, E.; Zilembo, N.; Ferrari, L.; Noberasco, C.;
Massaron, S.; Rimassa, L.; Bombardieri, E.:
Serum markers of bone metastases in postmenopausal breast cancer patients
treated with formestane.
Tumour Biol 18 (1997), 197-205
66. Mayne, P.D.; Thakar, S.; Rosalki, S.B.; Foo, A.Y.; Parbhoo, S.:
Identification of bone and liver metastases from breast cancer by measurement of
plasma alkaline phosphatase isoenzyme activity.
J Clin Pathol 40 (1987), 398-403
67. Mc Comb, R.B.; Bowers, G.N.; Posen, S.:
Alkaline Phosphatase.
Plenum Press, New York, (1979)

68. Miller, A.B.; Bulbrook, R.D.:
UICC multidisciplinary project on breast cancer: The epidemiology, aetiology and prevention of breast cancer.
Int J Cancer 37 (1986), 173-177
69. Möbus, V.; Kreienberg, R.:
Aktuelle Tumormarker beim Mammakarzinom.
In: Aktuelle Aspekte in der Diagnostik des Mammakarzinoms.
Hrsg: Jonat, W.
Bücherei des Frauenarztes 31 (1989), 62
70. Montravers, F.; Younsi, N.; Rousseau, C.; Philippe, C.; Seddiki, M.; Uzan, S.; Izrael, V.; Talbot, JN:
Comparison of bone scanning and CA 15-3 serum concentration in the follow-up of breast cancer.
Anticancer Res 17 (1997), 1683-1686
71. Murray, R.M.; Pitt, P.:
Medical adrenalectomy in patient with advanced breast cancer resistant to anti-oestrogen treatment.
Breast Cancer Res Treat 1 (1981), 91-95
72. Oremek, G.M.; Kramer, W.; Seiffert, U.B.; Jonas, D.:
Diagnostic value of skeletal AP and PSA with respect to skeletal scintigram in patients with prostatic disease.
Anticancer Res 17 (1997), 3035-3036
73. O'Mara, R.E.:
Skeletal scanning in neoplastic disease.
Cancer 37 (1976), 480-486
74. Ormiston, M. C.; Timoney, A. G.; Qureshi, A. R.:
Is follow up of patients after surgery for breast cancer worthwhile?
J R Soc Med 78 (1985), 920-921

75. Pandya, K.J.; Mc Fadden, E.T.; Kalish, L.A.; Tormey, D.C.; Taylor, S.G.; Falkson, G.:
A retrospective study of earliest indicators of recurrence in patients on Eastern Cooperative Oncology Group Adjuvant Chemotherapy Trials for breast cancer.
A preliminary report.
Cancer 55 (1985), 202-205
76. Panigrahi, K.; Delmas, P.D.; Singer, F.; Ryan, W.; Reiss, O.; Fisher, R.; Miller, P.D.; Mizrahi, I.; Darte, C.; Kress, B.C.:
Characteristics of a two-site immunoradiometric assay for human skeletal alkaline phosphatase in serum.
Clin Chem 40 (1994), 822-828
77. Paterson, C.R.; Robins, S.P.; Horobin, J.M.; Preece, P.E.; Cuschieri, A.:
Pyridinium crosslinks as markers of bone resorption in patients with breast cancer.
Br J Cancer 64 (1991), 884-886
78. Paulick, R.; Caffier, H.:
Klinische Bedeutung erhöhter CEA-Werte in der Nachsorge von Patientinnen mit Mammakarzinom.
Geburtsh Frauenheilk 45 (1985), 774-779
79. Paulick, R.; Caffier; H.; Paulick, M.:
Comparison of serum CEA, PHI, and TPA as tumor markers in breast cancer patients.
Cancer Detect Prev 10 (1987), 197-203
80. Pedrazzini, A.; Gelber, R.; Isley, M.; Castiglione, M.; Goldhirsch, A.:
First repeated bone scan in the observation of patients with operable breast cancer.
J Clin Oncol 4 (1986), 389-394

81. Perez, D.J.; Powles T.J.; Milan, J.; Gazet, J.C.; Ford, H.T.; Mc Cready, V.R.: Detection of breast carcinoma metastases in bone: relative merits of X-rays and skeletal scintigraphy.
Lancet 2 (1983), 613-616
82. Piovesan, A.; Berruti, A.; Torta, M.; Cannone, R.; Sperone, P.; Panero, A.; Gorzegno, G.; Termine, A.; Dogliotti, L.; Angeli, A.: Comparison of assay of total and bone-specific alkaline phosphatase in the assessment of osteoblast activity in patients with metastatic disease.
Calcif Tissue Int 61 (1997), 362-369
83. Plebani, M.; Bernadi, D.; Zaninotto, M.; de Paoli, M.; Secchiero, S.; Sciacovelli, L.: New and traditional serum markers of bone metabolism in the detection of skeletal metastases.
Clin Biochem 29 (1996), 67-72
84. Possinger,K.; Wilmanns,W.: Palliative Therapieführung zur Hemmung der Tumorprogression bei Patientinnen mit metastasierten Mammakarzinomen.
Internist (1993) 34: 340-350
85. Powels,T.J.: Advanced breast cancer-new approaches to treatment: a review.
Eur J Cancer Clin Oncol 21 (1988): 95-98
86. Preiss, J; Dornoff, W.; Hagmann, F.G.; Schmieder, A.: Mammakarzinom.
In: Empfehlungen zur Therapie.
Hrsg.: Onkologische Arbeitsgemeinschaft Saar-Pfalz-Mosel-e.V.
Onkologie 1998/99 (1998), 127-132

87. Reale, M.G.; Santini, D.; Marchei, G.G.; Manna, A.; Del Nero, A.; Marchei, P.: Skeletal alkaline phosphatase as a serum marker of bone metastases in the follow-up of patients with breast cancer.
Int J Biol Markers 10 (1995), 42-46
88. Rieden, K.:
Knochenmetastasen: Radiologische Diagnostik, Therapie und Nachsorge.
Springer-Verlag, Berlin, Heidelberg, New York, London, Tokyo, (1988)
89. Reitzig, P.:
Tumornachsorge beim Mammakarzinom-was ist sinnvoll.
Zeitschr Ärzt Fortbild 85 (1991), 1053
90. Roselli Del Turco, M.; Palli, D.; Cariddi, A.; Ciatto, S.; Pacini, P.; Distante, V.: Intensive diagnostic follow-up after treatment of primary breast cancer.
JAMA 271 (1994), 1593-1597
91. Rosen, P.; Groshen, W.; Saigo, P.; Kinne, D.W.; Hellman, S.:
A long-term follow-up study of survival in stage I (T1N0M0) and stage II (TiN1M0) breast carcinoma.
J Clin Oncol 7 (1989), 355-366
92. Rubens, R.D.; Fogelman, I.:
Bone metastases: diagnosis and treatment.
Springer, Berlin, Heidelberg, New York, London, Tokyo, (1991)
93. Rühl, U.:
Die Strahlentherapie nach Radikaloperation-Indikationen-Technik-Komplikationen.
Z ärztl Fortbild 85 (1991), 1045-1046
94. Rutgers, E.J.; van Slooten, E.A.; Kluck, H.M.:
Follow-up after treatment of primary breast cancer.
Br J Surg 76 (1989), 187-190

95. Scanlon, E.F.; Oviedo, M.A.; Cunningham, M.P.; Caprini, J.A.; Khandekar, J.D.: Preoperative and follow-up procedures on patients with breast cancer. Cancer 46 (1980), 977-979
96. Schapira, D.V.; Urban, N.: A minimalist policy for breast cancer surveillance. JAMA 265 (1991), 380-382
97. Schatz, C.; Lieven, H.V.; Mulders, M.; Rowold, J.; Stracke, H.; Müller, H.; Grebe, S.F.; Schatz, H.: Die Tumormarker CEA, TPA, und CA 19-9 sowie Ferritin und Osteocalcin in der Verlaufskontrolle beim Mammakarzinom. Wien Klin Wochenschr 23 (1985), 873-879
98. Schmid, L.; Schröck, R.; Langhammer, C.; Hünichen, A.; Bryxi, V.; Oberdorfer, M.: CA 15-3 und CEA in der Verlaufskontrolle des chemo-/hormontherapierten metastasierten Mammakarzinoms. In: Klinische Relevanz neuer monoklonaler Antikörper: 3. Hamburger Symposium über Tumormarker (1985). Hrsg: Greten, H. und Klapdor, R. Thieme, Stuttgart, (1986)
99. Schmidt-Rhode, P.; Rück, A.; Schulz, K. D.: Nachsorge des Mammakarzinoms. In: Gynäkologische Onkologie: Diagnostik, Therapie und Nachsorge bei Mamma- und Genital-Karzinomen in der Praxis. Hrsg: Mallmann, P. und Krebs, D. Zuckerschwerdt, München, Bern, Wien, New York, (1995), 195-205

100. Schreiber, R.R.:
The radiologist and the diagnosis of bone metastasis.
In: Bone Metastasis.
Hrsg: Weiss, L.; Gilbert, H.A.; Hall, G.K.
Medical publishers, Boston, (1981)
101. Schünemann, H.; Jourdain, M.:
Stellenwert der Skelettszintigraphie in der Nachsorge des Mammakarzinoms.
Deutsche Med Wochenschrift 114 (1989), 3-8
102. Schumacher, T.; Schmitz, S.; Diehl, V.:
Prognoserelevanz der onkologischen Nachsorge.
Versicherungsmedizin 47 (1995), 166-170
103. Seifert, M.; Adler, A.; Obermair, A.; Hartmann, B.; Denison, U.; Kubista, E.;
Sevelda, P.:
Möglichkeiten und Ziele in der Tumornachsorge beim Mammakarzinom.
Gynakol Rundsch 31 (1991), 309-311
104. Sekine, H.; Hayes, D.F.; Ohno, T.; Keefe, K.A.; Schaetzl, E.; Bast, R.C.;
Knapp, R.:
Circulating DF3 and CA 12-5 antigen levels in serum from patients with epithelial
ovarian carcinoma.
J Clin Oncol 3 (1985), 1355-1363
105. Shinozaki, T.; Chigira, M.; Kato, K.:
Multivariate analysis of serum tumor markers for diagnosis of skeletal
metastases.
Cancer 69 (1992), 108-112

106. Spiessl, B.; Beahrs, O.H.; Hermanek, P.; Hutter, R.V.P.; Scheibe, O.; Sabin, L.H.; Wagner, G.:
Mammatumoren.
In: TNM-Atlas. Illustrierter Leitfaden zur TNM/pTNM-Klassifikation maligner Tumoren.
Springer, Berlin, Heidelberg, New York, London, Tokyo, (1993)
- 107.. Stegner, H.E.:
Gynäkologie und Geburtshilfe.
Enke, (1996)
108. Stieber, P.; Nagel, D.; Ritzke, C.; Rössler, N.; Kirsch, C.M.; Eiermann, W.; Fateh-Moghadam, A.:
Significance of bone alkaline phosphatase, CA 15-3 and CEA in the detection of bone metastases during the follow-up of patients suffering from breast carcinoma.
Eur J Clin Chem Clin Biochem 30 (1992), 809-814
109. Smalley, R.V.; Scogna, D.M.; Malmud, L.S.:
Advanced breast cancer with bone-only metastases. A chemotherapeutically responsive pattern of metastases.
Am J Clin Oncol 5 (1982) 161-166
110. The GIVIO Investigators:
Impact of follow-up and testing on survival and health-related quality of life in breast cancer patients: a multicenter randomized trial.
JAMA 272 (1994), 1587-1592
111. Theriault, R.L.; Hortobagyi, G.N.:
Bone metastases in breast cancer.
Anticancer Drugs 3 (1992), 455-462

112. Tobias, R.; Rithwell, C.; Wagner, J.; Green, A.; Liu, Y.S.V.:
Development and evaluation of a radioimmunoassay for the detection of a
monoclonal antibody defined breast tumor associated antigen 115D8/DF3.
Kongress der American Association of Clinical Chemistry (AACC), (Atlanta 1985)
113. Tomin, R.; Donegan, W.:
Screening for recurrent breast cancer - its effectiveness and prognostic value.
J Clin Oncol 5 (1987), 62-67
114. Tondini, C.; Hayes, D.F.; Gelmann, R.; Henderson, I.C.; Kufe, D.W.:
Comparison of CA 15-3 and CEA in monitoring the clinical course of patients with
metastatic breast cancer.
Cancer Res 48 (1988), 4107-4112
115. Umbach, G.E.; Holzki, C.; Perschman, U.; Schnurch, H.G.; Bender, H.G.:
Erfahrungen in der Nachsorge von Patientinnen mit Mammakarzinom.
Geburtsh Frauenheilk 48 (1988), 292-298
116. van Dalen, A.:
TPS in breast cancer-a comparative study with carcinoembryonic antigen and CA
15-3.
Tumour Biol 13 (1992), 10-17
117. van Holten-Verzantvoort, A.T.; Bijvoet Olm Hermans, J.; Cleton, F.J.; Krohn,
H.M.; Harinck, H.I.; Vermey, P.; Elite, J.W.; Neyt, J.P.; Beex, L.V.:
Reduced morbidity from skeletal metastases in breast cancer patients during
long-term bisphosphonate (APD) treatment.
Lancet 2 (1987), 983-985
118. Wandt, H.; Bruntsch, U.; Gallmeier, W.M.:
Nachsorge beim Mammakarzinom.
Deutsche Med Wochenschrift 114 (1989), 1130-1136

119. Webber, W.; Boyd, N.:
A critique of the methodology of studies of benign breast disease and breast cancer risk.
J Natl Cancer Inst 77 (1986), 397-404
120. Weiss, M.; Cole, D.; Ray, K.; Whyte, M.P.; Lafferty, M.A.; Mulivor, R.A.; Harris, H.:
A missense mutation in the human liver-bone-kidney alkaline phosphatase gene causing a lethal form of hypophosphatasia.
Proc Natl Acad Scie U S A 85 (1988), 7666-7669
121. Wertheimer, M.D.:
Against minimalism in breast cancer follow-up.
JAMA 265 (1991), 396-397
122. Westerhuis, L. W.; Delaere, K. P.:
Diagnostic value of some biochemical bone markers for the detection of bone metastases in prostate cancer.
Eur J Clin Chem Clin Biochem 35 (1997), 89-94
123. Winchester, D.P.; Sener; S.F., Khandekar, J. D.; Oviedo, M.A.; Cunningham, M.P.; Caprini, J.A., Burkett, F.E., Scanlon, E.F.:
Symptomatology as an indicator of recurrent or metastatic breast cancer.
Cancer 43 (1979), 956-960
124. Winzer, J.W.:
Die chirurgische Therapie beim nichtinvasiven Karzinom der Mamma.
Z ärztl Fortbild 85 (1991), 1041-1044
125. Withold, W.; Gorgescu, G.; Khakzad, H.; Vosberg, H.; Mueller-Gaertner, H.W.; Reinauer, H.:
Efficiency of simultaneous determination of bone alkaline phosphatase mass concentration in serum and urinary excretion of pyridinium cross-links for detection of bone metastases.
Clin Biochem 28 (1995), 511-517

126. Wolff, J. M.; Ittel, T.; Boeckmann, W.; Reinike, T.; Habib, F. K.; Jaske, G.:
Skeletal alkaline phosphatase in the metastatic workup of patients with prostate
cancer.
Eur Urol 30 (1996), 302-306
127. Yamashita, K.; Ueda, T.; Takafumi, U.; Komatsubara, Y.; Koyama, H., Inaji, H.;
Yonenobu, K.; Ono, K.:
Breast cancer with bone-only metastases. Visceral metastases-free rate in
relation to anatomic distribution of bone metastases.
Cancer 68 (1991), 634-637
128. Zanco, P.; Rota, G.; Sportiello, V.; Borsato, N.; Ferlin, G.:
Diagnosis of bone and liver metastases in breast cancer comparing tumor
markers and imaging techniques.
Int J Biol Markers 4 (1989), 103-105
129. Zaninotto, M.; Secchiero, S.; Rubin, D.; Sciacovelli, L.; Trovo, M.; Bortolus, R.;
Plebani, M.:
Serum bone alkaline phosphatase in the follow-up of skeletal metastases.
Anticancer Res 15 (1995), 2223-2228
130. Zwaveling, A.; Albers, G.H.; Felthuis, W.; Cronin, W.:
An evaluation of routine follow-up for detection of breast cancer recurrences.
J Surg Oncol 34 (1987), 194-197