

6. Literatur

1. Burgdorfer W, Barbour AG. Lyme disease - a tick-borne spirochetosis? *Science* 1982;261:1317-1319
2. Steere AC. Lyme disease. *N. Engl. J. Med.* 1989;321:586-597
3. Johnson RC, Schmid GP, Hyde FW, Steigerwalt AG, Brenner DJ. *Borrelia burgdorferi* sp. nov.: etiologic agent of Lyme disease. *Int. J. Syst. Bact.* 1984;34:496-497
4. Johnson RC, Hyde FW, Rumpel CM. Taxonomy of the Lyme disease spirochetes. *Yale J. Biol. Med.* 1984;57:529-537
5. Canale-Parola E. Order spirochaetales. In: Kiege NR, Holt JG, eds. *Bergey's Manual of Systemic Bacteriology*. Vol. Vol. 1. Baltimore: Williams & Williams:38-39
6. Karlsson M. Comparison of western blot and enzyme-linked immunosorbent assay for diagnosis of Lyme borreliosis. *Eur. J. Clin. Microbiol.* 1989;8:871-877
7. Ma B. Serodiagnosis of lyme borreliosis by western immunoblot: reactivity of various significant antibodies against *B. burgdorferi*. *J. Clin. Microbiol.* 1992;30:370-376
8. Magnarelli LA, Anderson JF. Enzyme-linked immunoabsorbent assays for the detection of class-specific immunoglobulins to *B. burgdorferi*. *Am. J. Epidemiol.* 1988;127:813-815
9. Russell H, Sampson JS, Schmidt GP. Enzyme-linked immunoabsorbent assays and indirect immunofluorescence assay for Lyme disease. *J. Infect. Dis.* 1984;149:465-470
10. Norris SJ, Carter JC, Howell KJ, Barbour AG. Low-passage-associated proteins of *Borrelia burgdorferi* B31: characterization and molecular cloning of Osp D, a surface exposed, plasmid-encoded lipoprotein. *Infect. Immun.* 1992;60:4662-4672
11. Dressler F, Steere AC. Western blotting in the serodiagnostic of Lyme disease. *J. Infect. Dis.* 1993;167:392-400
12. Brandt ME, Riley BS, Radolf JD, Norgard MV. Immunogenic integral membrane proteins of *Borrelia burgdorferi* are lipoproteins. *Infect. Immun.* 1990;58:983-991
13. Chamberlain NR, Brandt ME, Erwin AL, Radolf JD, Norgard MV. Major integral membrane protein immunogens of *Treponema pallidum* are proteolipids. *Infect. Immun.* 1989;57:2872-2877
14. Schouls LM, Mout R, Dekker J, Emden JDAv. Characterization of lipid-modified immunogenic lipids of *Treponema pallidum* expressed in *Escherichia coli*. *Microbiol. Pathol.* 1989;7:175-188

15. Swancutt MA, Radolf JD, Norgard MV. The 34-kD membrane immunogen of *Treponema pallidum* is a lipoprotein. *Infect. Immun.* 1990;58:384-392
16. Bergstrom S, Barbour AG, Bundoc VG. Molecular analysis of linear plasmid-encoded major surface proteins, Osp A and Osp B of the Lyme disease spirochete *Borrelia burgdorferi*. *Mol. Microbiol.* 1989;3:479-486
17. Fuchs R, Jauris F, Lottspeich F, Praec-Mursic V, Wilske B, Soutschek E. Molecular analysis and expression of a *Borrelia burgdorferi* gene encoding a 22 kDa protein (pC) in *Escherichia coli*. *Mol. Microbiol.* 1992;6:503-509
18. Lam TT, Nguyen TPK, Montgomery RR, Kantor FS, Fikrig E, Flavell RA. Outer surface proteins E and F of *Borrelia burgdorferi*, the agent of Lyme disease. *Infect. Immun.* 1994;62:290-298
19. Wilske B, Praec-Mursic S, Jauris A, Hofmann A, Pradel I, Soutschek E. Immunological and molecular polymorphisms of Osp C, and immunodominant major outer surface protein of *Borrelia burgdorferi*. *Infect. Immun.* 1993;61:2182-2191
20. Akins DR, Purcell BK, Mitra M, Norgard MV, Radolf JD. Lipid modification of the 17-kD membrane immunogen of *Treponema pallidum* determines macrophage activation as well as amphiphilicity. *Infect. Immun.* 1993;61:1202-1210
21. Chamberlain NR, DeOgny L, Slaughter C, Radolf JD, Norgard MV. Acylation of the 47-kD major membrane immunogen of *Treponema pallidum* determines its hydrophobicity. *Infect. Immun.* 1989;57:2878-2885
22. Purcell BK, Swancutt MA, Radolf JD. Lipid modification of the 15-kD major membrane immunogen of *Treponema pallidum*. *Molec. Microbiol.* 1990;4:1371-1379
23. Weigel LM, Brandt ME, Norgard MV. Analysis of the N-terminal region of the 47-kD integral membrane lipoprotein of *Treponema pallidum*. *Infect. Immun.* 1992;60:1568-1576
24. Ma Y, Weis JJ. *Borrelia burgdorferi* outer surface lipoproteins Osp A and Osp B possess B-cell mitogenic and cytokine-stimulatory properties. *Infect. Immun.* 1993;61:3843-3853
25. Radolf JD, Norgard MV, Brandt ME, Isaacs RD, Thompson PA, Beutker B. Lipoproteins of *Borrelia burgdorferi* and *Treponema pallidum* activate cachectin/tumor necrosis factor synthesis: analysis using a CAT reporter construct. *J. Immunol.* 1991;147:1968-1974

26. Riley BS, Oppenheimer-Marks N, Hansen EJ, Radolf JD, Norgard MV. Virulent *Treponema pallidum* activates human vascular endothelial cells. *J. Infect. Dis.* 1992;165:484-493
27. Erdile LF, Brandt MA, Warakomski DJ, Westrack GJ, Sadziene A, Barbour AG, Mayes JP. Role of attached lipid in immunogenicity of *Borrelia burgdorferi* Osp A. *Infect. Immun.* 1993;61:81-90
28. Deres K, Schild H, Wiesmüller KH, Jung G, Rammensee HG. In vivo priming of virus-specific cytotoxic T lymphocytes with synthetic lipopeptide vaccine. *Nature* 1989;342:561-564
29. Hausschildt S, Hoffman P, Beuscher HU, Dufhues G, Heinrich P, Wiesmüller KH. Activation of bone marrow-derived mouse macrophages by bacterial lipopeptide: cytokine production, phagocytosis, and Ia expression. *Eur. J. Immunol.* 1990;20:63-68
30. Metzger R, Jung G, Bessler WG, Hoffmann P, Strecker M, Lieberknecht A, Schmidt U. Lipopeptides containing 2-(palmitoylamino)-6,7-bis(palmitoxy)heptanoic acid: synthesis, stereospecific stimulation of B lymphocytes and macrophages, and adjuvant activity in vivo and in vitro. *J. Med. Chem.* 1991;34:1969-1974
31. Aguero-Rosenfeld MA. Evolution of the serologic response to *Borrelia burgdorferi* in treated patients treated with culture-confirmed erythema migrans. *J. Clin. Microbiol.* 1996;34:1-9
32. Wilske B. Antigenic variation of *Borrelia burgdorferi*. *Ann. N.Y. Acad. Sci.* 1988;26:126-143
33. Marconi RT, Garon CF. Variability of osp genes and gene products among species of Lyme disease spirochetes. *Infect. Immun.* 1993;61:2611-2617
34. Schwan TG, Warren GS. Factors influencing the antigenic reactivity of *Borrelia burgdorferi*, the Lyme disease spirochete. *Scand. J. Infect. Dis.* 1991;77:94-101
35. Cinco M, Banfi E, Panfili E. Leptospiral lipopolysaccharide presence in the outer envelope: electrophoretic and immunological specificity. *Zbl. Bakt. Hyg.* 1988;269:277-283
36. Adler B, Ballard SA, Miller SJ, Faine S. Monoclonal antibodies reacting with serogroup and serovar specific epitopes on different lipopolysaccharide subunits of *Leptospira interrogans* serovar pomona. *FEMS Microbiol. Immunol.* 1989;47:213-218

37. Vinh T, Shi M, Adler B, Faine S. Characterization and taxonomic significance of lipopolysaccharides of *Leptospira interrogans* serovar hardjo. J. Gen. Microbiol. 1989;135:2663-2673
38. Halter MR, Joens LA. Lipopolysaccharides from *Treponema hyodysenteriae* and *Treponema innocens*. Infect. Immun. 1988;56:3152-3156
39. Takayama K, Rothenberg RJ, Babour AG. Absence of lipopolysaccharide in the Lyme disease spirochete, *Borrelia burgdorferi*. Infect. Immun. 1987;55:2311-2313
40. Cinco M, Banfi E, Balanzin D, Godeas C, Panfili E. Evidence for (lipo)oligosaccharides in *Borrelia burgdorferi* and their serological specificity. FEMS Microbiol. Immunol. 1991;76:33-38
41. Galanos C, Lüderitz O, Westphal O. A new method for the extraction of R lipopolysaccharides. Eur. J. Biochem. 1969;9:245-249
42. Habicht GS, Beck G, Coleman JL. *Borrelia burgdorferi* lipopolysaccharides and its role in the pathogenesis of Lyme disease. Zbl. Bakt. Mikrobiol. Hyg. 1985;263:137-141
43. Tsai CM, Frasch CA. A sensitive silver stain for detecting lipopolysaccharides in polyacrylamide gels. Anal. Biochem. 1982;119:115-119
44. Eiffert H, Lotter H, Jarecki-Khan K, Thompsen R. Identification of an immunoreactive non-proteinaceous component in *Borrelia burgdorferi*. Med. Microbiol. Immunol. 1991;180:229-237
45. Wheeler CM, Monco JCG, Benach JL, Golightly MG, Habicht GS, Steere AC. Nonprotein antigens of *Borrelia burgdorferi*. J. Infect. Dis. 1993;167:665-674
46. Beck G, Habicht GS, Benach JL, Coleman JL. Chemical and biological characterization of a lipopolysaccharide extracted from the Lyme disease spirochete (*Borrelia burgdorferi*). J. Infect. Dis. 1985;152:108-117
47. Radolf JD, Goldberg MS, Bourell K, Baker SI, Jones JD, Norgard MV. Characterization of outer membranes isolated from *Borrelia burgdorferi*, the Lyme disease spirochete. Infect. Immun. 1995;63:2154-2163
48. Raoult D, Hechemy E, Baranton G. Cross-reaction with *Borrelia burgdorferi* antigen of sera from patients with human immunodeficiency virus infection, syphilis and leptospirosis. J. Clin. Microbiol. 1989;27:2152-2155
49. Livermore BP. Lipids of the spirochetes: Comparison of the lipids of several members of the genera *Spirocheta*, *Treponema* and *Leptospira*. J. Bact. 1974;120:1268-1273

-
50. Herbert MM, Tze-Ken Y, Jenkin HM. Alk-1-enyl ether phospholipids (plasmalogens) and glycolipids of *Treponema hyodysenteriae*. *Biochim. Biophys. Acta* 1980;618:273-281
 51. Livermoore BP, Johnson RC. Isolation and characterization of a glycolipid from *Treponema pallidum*, Kazan 5. *Biochim. Biophys. Acta* 1970;210:315-318
 52. Radolf JD, Robinson EJ, Bourell KW. Characterization of outer membranes isolated from *Treponema pallidum*, the syphilis spirochete. *Infect. Immun.* 1995;63:4244-4252
 53. Pickett J, Kelly R. Lipid catabolism of relapsing fever borreliae. *Infect. Immun.* 1974;9:279-285
 54. Livermoore BP, Bey RF, Johnson RC. Lipid metabolism of *Borrelia hermsii*. *Infect. Immun.* 1978;20:215-220
 55. Cacciapuoti B, Ciceroni L, Barbini DA. Fatty acid profiles in the family Leptospiraceae. *Zbl. Bakt.* 1990;274:16-27
 56. Cho SN, Uhm JR, Kim JD. Comparative analysis of lipopolysaccharide and lipid antigens of *Leptospira interrogans* serovars. *Yonsei Med. J.* 1992;33:24-31
 57. Ono E, Takase H, Naiki M, Yanagawa R. Purification, characterization and serological properties of a glycolipid antigen reactive with a serovar specific monoclonal antibody against *Leptospira interrogans* serovar canicola. *J. Gen. Microbiol.* 1987;133:1329-1336
 58. Masuzawa T, Nakamura R, Shimizu T, Iwamoto Y, Morita T, Yanagihara Y. Immunological characteristics of the glycolipid antigen of *Leptospira interrogans* serovar lai. *Infect. Immun.* 1989;57:2502-2506
 59. Moribayashi A, Goto N, Arimitsu Y, Himeno K, Tatsuki S. Lipids and fatty acids of *Leptospira interrogans* serovar copenhageni virulent strain shibaura. *Jpn. J. Sci. Biol.* 1991;44:87-97
 60. Barbour AG. Isolation and cultivation of Lyme Disease spirochetes. *Yale J. Biol. Med.* 1984;57:521-525
 61. Kelly R. Cultivation of *Borrelia hermsii*. *Science* 1971;173:443-444
 62. Stoenner HG. Biology of *Borrelia hermsii* in Kelly medium. *Appl. Microbiol.* 1974;28:540-543
 63. Kates M. Techniques in lipidology. Isolation, analysis and identification of lipids. In: Work TS, Work E, eds. *Laboratory techniques in biochemistry and molecular biology*. Vol. 3. Amsterdam: North-Holland Publishing Co., 1972:269-610

-
64. Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* 1959;37:911-917
 65. Williams MA, McCluer RH. The use of Sep-Pak C18 cartridges during the isolation of gangliosides. *J. Neurochem.* 1980;35:266-269
 66. Kaluzny MA, Duncan LA, Merritt MV, Epps DE. Rapid separation of lipid classes in high yield and purity using bonded phase columns. *J. Lipid Res.* 1985;26:135-140
 67. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680-685
 68. Towbin H, Staehlin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA* 1979;76:4350-4354
 69. Hensel U, Wellensiek HJ, Bhakdi S. Sodium dodecyl sulfate polyacrylamide gel electrophoresis immunoblotting as a serological test in the diagnosis of syphilic infections. *Am. Soc. of Microbiol.* 1985;21:82-87
 70. Alfen I, Wellensiek HJ. Die Bedeutung kreuzreagierender Antikörper für die Serodiagnostik der Lyme-Borreliose und der Syphilis. *Lab. Med.* 1994;18:12-19
 71. Dittmer JC, Lester RL. A simple, specific spray for the detection of phospholipids on thin-layer chromatograms. *J. Lipid Res.* 1964;5:126-127
 72. Baumeister S, Dennis RD, Klünder R, Schares G, Zahner H, Geyer E. Litomosoides carinii: macrofilariae-derived glycolipids - chromatography, serology and potential in the evaluation of anthelmintic efficacy. *Parasite Immunol.* 1994;16:629-641
 73. Geyer R, Geyer H, Kühnhardt S, Mink W, Stirm S. Capillary gas-chromatography of methylhexitol acetates obtained upon methylation of N-glycosidically linked glycoprotein oligosaccharides. *Anal. Biochem.* 1982;121:263-274
 74. Paz-Parente J, Cardon P, Leroy Y, Montreuil J, Fournet B, Ricard G. A convenient method for methylation of glycoprotein glycans in small amounts by using lithium methyl-sulfinyl carbanion. *Carbohydr. Res.* 1985;141:41-47
 75. Geyer R, Geyer H. Saccharide linkage analysis using methylation and other techniques. *Methods Enzymol.* 1994;230:86-108
 76. Dell A. Preparation and desorption mass spectrometry of permethyl and peracetyl derivatives of oligosaccharides. *Methods Enzymol.* 1990;193:647-660

-
77. Hoffman J, Lindberg B, Svensson S. Determination of the anomeric configuration of sugar residues in acetylated oligo- and polysaccharides by oxidation with chromium trioxide in acetic acid. *Acta Chem. Scand.* 1972;26:661-666
 78. Gaver RC, Sweeley CC. Methods for methanolysis of sphingolipids and direct determination of long-chain bases by gas chromatography. *J. Am. Oil Chem. Soc.* 1965;42:294-298
 79. Livermore BP, Johnson RC. Lipids of the Spirochaetales: comparison of the lipids of several members of the genera Spirochaeta, Treponema and Leptospira. *J. Bacteriol.* 1974;120:1268-1273
 80. Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb JF, Fleischmann RD, Richardson D, Peterson J, Kerlavage AR, Quackenbush J, Salzberg S, Hanson M, Vugt Rv, Palmer N, Adams MD, Gocayne J, Weidman J, Utterback T, Wathley L, McDonald L, Artiach P, Bowman C, Garland S, Fujii C, Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter JC. Genomic sequence of a lyme disease spirochaete, *Borrelia burgdorferi*. *Nature* 1997;390:580-586
 81. Livermore BP, Johnson RC. Isolation and characterization of a glycolipid from *Treponema pallidum*, Kazan 5. *Biochem. Biophys. Acta* 1970; 210; 315-318
 82. Kawabata H, Masuzawa T, Yanagihara Y. Genomic analysis of *Borrelia japonica* sp. nov. isolated from Ixodes ovatus in Japan. *Microbiol Immunol.* 1993; 37; 843-848
 83. Marconi RT, Liveris D, Schwartz I. Identification of novel insertion elements, restriction fragment length polymorphisms patterns, and discontinuous 23S rRNA in Lyme Disease Spirochetes: Phylogenetic analyses of rRNA genes and their intergenic spacers in *Borrelia japonica* sp. nov. and genomic group 21038 (*Borrelia andersonii* sp. nov.) isolates. *J. Clin. Microbiol.* 1995; 33; 2427-2434
 84. Wang G, Van Dam AP, Le Fleche A, Postic D, Peter O, Baranton G, De Boer R, Spanjaard L, Dankert J. Genetic and phenotypic analysis of *Borrelia valaisiana* sp. nov. (*Borrelia* genomic group VS116 and M19)
 85. Assous MV, Postic D, Paul G, Nevot P, Baranton G. Individualisation of two new genomic groups among American *Borrelia burgdorferi* sensu lato strains. *FEMS Microbiol Lett.* 1994; 121; 93-98
 86. Postic D, Marti Ras N, Lane RS, Henderson M, Baranton G. Expanded diversity among Californian *Borrelia* isolates and description of *Borrelia bissettii* sp. nov. (formerly *Borrelia* group DN 127)

87. Singer SJ, Nicolson GL. The fluid mosaic model of the structure of cell membranes. *Science* 1972; 175; 720-731
88. Finean J, Coleman R, Mitchell RH. *Membranes and their cellular functions*. 3rd Edt. Blackwell Scientific Publications 1984, Oxford
89. Rothman JE, Lenard J. Membrane asymetry. *Science* 1977; 195; 743-753
90. Vance DE, Vance JE. *Biochemistry of lipids, Lipoproteins and membranes*. New Comprehensive Biochemistry 1991, Vol. 20, Elsevier Science Publishing Co., Inc. NY
91. Wiegandt H. Glycosphingolipids. *Adv. Lipid Res.* 1971; 9, 249-289
92. Hakomori SI. Glycosphingolipide. *Spektrum der Wissenschaft* (Juli); 90-100
93. Sastry PS. Lipids of nervous tissue: Composition and metabolism. *Prog. Lipid Res.* 1985; 24; 169-176
94. Nagai Y. Gangliosides as signalling molecules in neural functions. *Glycoconj. J.* 1995; 12; 544
95. Ishizuka I Yamakawa T. Glycoglycerolipids. *Glycolipids* (H. Wiegandt, Ed.) 1985; 101-198; Elsevier, Amsterdam
96. Schwarting GA, Gajewski A. Glycolipids of murine lymphocyte subpopulations: a defect in the levels of sialidase-sensitive sialosylated asialo GM1 in beige mouse lymphocytes. *J. Immunol.* 1981; 126; 2403-2407
97. Hakomori SI. Bloodgroup ABH and Ii antigens of human erythrocytes: chemistry, polymorphism, and their, developmental change. *Sem. Haematol.* 1981, 18; 39-62
98. Watkins WM. Biosynthesis: molecular basis of antigenic specificity in the ABO, H and Lewis blood group system. *Glycoproteins* (J. Montreuil, H. Schachter, J.F.G. Vliegthart, Ed.) 1995; 313-390
99. Vanier MT. Lipid changes in Niemann-Pick disease type C brain: personal experience and review of the literature. *Neurochem Res.* 1999 Review; 24(4); 481-489.
100. Geyer R, Geyer H, Kühnhardt S, Mink W Stirm S. Methylation analysis of complex carbohydrates in small amounts: Capillary gas chromatography – mass fragmentography of methylalдитiol acetates obtained from N-glycosidically linked glycoprotein oligosaccharides. *Anal. Biochem.* 1983; 133; 197-207
101. Afzelius A. Erythema chronicum migrans. *Acta Derm. Venereol.* (Stockholm) 1921; 2;120-5

102. Rille JH. Acrodermatitis chronica atrophicans. Arch. Dermatol. Syph. (Berlin) 1898; 45;423
103. Herxheimer K, Hartmann K. Ueber acrodermatitis chronica atrophicans. Arch. Dermatol. Syph. 61; 57-76
104. Lipschütz B. Ueber eine seltene Erythemform (*Erythema chronicum migrans*). Arch. Dermatol. Syph. 1914; 118; 349-356.
105. Hellerström S. Beitrag zur Pathogenese des Erythema chronicum migrans Afzelii. Acta Derm Venereol. (Stockh) 1934; 14; 517
106. Garin CH, Bujadoux CH. Paralyse par les tiques. J. Med. Lyon 1922; 71; 765-767
107. Hellerström S. Erythema chronicum migrans Afzelii. Acta Derm Venereol. (Stockh) 1930; 11; 315-321
108. Bannwarth A. Zur Klinik und Pathogenese der chronische lymphozytären Meningitis. Arch. Psych. 1944; 117; 161-185 und 682-717
109. Hörstrup P, Ackermann R. Durch Zecken übertragene Meningopolyneuritis (Garin-Bujadoux-Bannwarth). Fortschr. Neurol. Psych. 1973; 41;583-606
110. Bäfverstedt B. Ueber Lymphadenosis benigna cutis. Eine klinische und pathologisch-anatomische Studie. Acta. Derm. Venereol. (Stockh) 1943; 24; suppl. 11; 1-102
111. Jordan P, Holtschmidt J. Traumatisches Zeckenbiß-Lymphozytom und Erythema chronicum migrans. Hautarzt 1951; 2; 397-401
112. Bianchi E. Die Penicillinbehandlung der Lymphozytome. Dermatologica 1950; 100; 270-273
113. Paschoud JM. Die Lymphadenosis benigna cutis als übertragbare Infektionskrankheit I-III. Hautarzt 1958; 9; 153-165, 263-269 und 311-315
114. Steere AC, Malawista SE, Snyderman DR, Shope RE, Andiman WA, Ross MR, Steele FM. Lyme Arthritis: An epidemic of oligoarticular arthritis in children and adults in three Connecticut communities. Arthr. Rheum. 1977; 20: 7-17
115. Barbour AG. Isolation and cultivation of Lyme disease spirochetes. Yale J. Biol. Med. 1984; 57; 521-525
116. Benach JL, Bosler EM, Hanrahan JP, Coleman JL, Habicht JS, Bast TF, Cameron DJ, Ziegler JL, Barbour AG, Burgdorfer W, Edelmann R, Kaslow RA. Spirochetes isolated from blood of two patients with Lyme disease. New Engl. J. Med. 1983; 308; 740-742
117. Holt, SC. Anatomy and chemistry of spirochetes. Microbiol. Rev. 1978;42; 144-160

118. Hovind-Hougen, K. Ultrastructure of spirochetes isolated from *Ixodes ricinus* and *Ixodes dammini*. Yale J. Biol. 1984; 57; 543-548
119. Barbour AG, Hayes SF. Biology of *Borrelia* species. Microbiol. Rev. 1986; 50; 381-400
120. Barbour AG, Hayes SF, Heiland RA, Schrupf ME, Tessier SL. A borrelia-specific monoclonal antibody binds to a flagellar epitope. Infect. Immun. 1986; 52; 549-554
121. Adam T, Gassmann GS, Rasiah C, Göbel UB. Phenotypic and genotypic analysis of *Borrelia burgdorferi* isolates from various sources. Infect. Immun. 1991; 59; 2579-2585
122. Baranton G, Portic D, Girons JS. Delineation of *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* sp. Novii, and group VS461 associated with Lyme borreliosis. Int. J. syst. Bacteriol. 1992; 42; 378-383
123. Belfaiza J, Postic D, Bellenger E, Baranton G, Girons JS. Genomic fingerprinting of *Borrelia burgdorferi* sensu lato by pulsed-field gel electrophoresis. J. Clin. Microbiol. 1993; 31; 2873-2877
124. Canica MM, Nato F, Du Merle L, Mazie JC, Baranton G, Postic D. Monoclonal antibodies for identification of *Borrelia afzelii* sp. Nov. Associated with late cutaneous manifestations of Lyme borreliosis. J. Infect. Dis. 1993; 25; 441-448
125. Wallich R; Helmes C, Schaible UE, Lobet Y, Moter SE, Kramer MD, Simon MM. Evaluation of genetic divergence among *Borrelia burgdorferi* isolates by use of Osp A, fla, HSP 60 and HSP 70 probes. Infect. Immun. 1992; 60; 4856-4866
126. Callister SM, Agger WA, Schell RF, Johnson RC, Ellingson JLE. Effects of bovine serum albumin on the ability of Barbour-Stoenner-kelly Medium to detect *Borrelia burgdorferi*. J. Clin. Microbiol. 1990; 28; 363-365
127. Bergmann J, Liebisch A, Pohlmeier K. Borreliose – zum Vorkommen der einheimischen Borreliose bei Zecken, Wild- und Haustieren in einem niedersächsischen Moor. J. für den Veterinär 1992; 2; 12-15
128. Aeschlimann A, Chamot E, Gigon F, Jeanneret JP, Kessler D, Walther C. *Borrelia burgdorferi* in Switzerland. Zentralbl. Bakt. Hyg. 1986; A263; 450-458
129. Anderson JF. Epizootiology of *Borrelia* in *Ixodes* tick vectors and reservoir hosts. Rev. Infect. Dis. 1989; 11, suppl. 6, 1451-1458
130. Walter G, Liebisch A. Untersuchungen zur Biologie und Verbreitung von Zecken in Norddeutschland. Z. Angew. Zool. 1980; 67; 449-476
131. Kurtenbach K, Dizij A, Seitz HM, Margos G, Moter SE, Kramer MD, Wallich R, Schaible UE, Simon MM. Differential immune responses to *Borrelia burgdorferi* in

- European wild rodent species influence spirochete transmission to *Ixodes ricinus*. Infect. Immun. 1994; 62; 5344-5352
132. Liebisch A, Schmidt C, Friedhoff KT. Zeckenbefall und durch Zecken übertragene Infektionen bei kleinen Haustieren unter Berücksichtigung der Lyme Borreliose. Der praktische Tierarzt 1993; 23; 112-115
133. Liebisch A. Biologie und Ökologie der Zecken. Horst H. (Hrsg.): Einheimische Zeckenborreliose (Lyme-Krankheit) bei Mensch und Tier. Perimed Erlangen 1991; 31-47
134. Burgdorfer W, Hayes SF, Corwin D. Pathophysiology of the Lyme disease spirochete, *Borrelia burgdorferi* in Ixodes ticks. Rev. Infect. Dis. 1989; 11; suppl. 6; 1442-1450
135. Gern L, Aeschlimann A. Development of *Borrelia burgdorferi* in *Ixodes ricinus* during blood meal. IV International Conference on Lyme Borreliosis, Stockholm 1990 (Abstr.)
136. Piesman J, Oliver JR, Sinsky R. Growth kinetics of the Lyme disease spirochete *Borrelia burgdorferi* in vector ticks (*Ixodes dammini*). Am. J. Trop. Med. Hyg. 1990; 42; 352-357
137. Piesman J, Mather TN, Sinsky R, Spielman A. Duration of tick attachment and *Borrelia burgdorferi* transmission. J. Clin. Microbiol. 1987; 25; 557-558
138. Kaneda K, Masuzawa T, Yasugami K, Suzuki T, Suzuki Y, Yanagihara Y. Glycosphingolipid-binding protein of *Borrelia burgdorferi* sensu lato. Infect. Immun. 1997; 65 (8); 3180-3185
139. Backenson PB, Coleman JL, Benach JL. *Borrelia burgdorferi* shows specificity of binding to glycosphingolipids. Infect. Immun. 1995; 63 (8); 2811-2817
140. Norgard MV, Riley BS, Richardson JA, Radolf JD. Dermal inflammation elicited by sythetic analogs of *Treponema pallidum* and *Borrelia burgdorferi* lipoproteins. Infect. Immun. 1995; 63 (4); 1507-1515
141. Skare JT, Mirzabekov TA, Shang ES, Blanco DR, Erdjument-Bromage H, Bunikis H, Bergstrom S, Tempst P, Kagan BL, Miller JN, Lovett MA. The Oms66 (p66) protein is a *Borrelia burgdorferi* porin. Infect. Immun. 1997; 65 (9); 3654-3661
142. Belisle JT, Brandt ME, Radolf JD, Norgard MV. Fatty acids of *Treponema pallidum* and *B. burgdorferi* lipoproteins. J. Bacteriol. 1994; 176 (8); 2151-2157
143. Garcia-Monco JC, Fernandez Villar B, Rogers RC, Szczepanski A, Wheeler CM, Benach JL. *Borrelia burgdorferi* and other related spirochetes bind to galactocerebroside. Neurology 1992; 42 (7); 1341-1348

144. Katona LI, Beck G, Habicht GS. Purification and immunological characterization of a major low-molecular-weight lipoprotein from *Borrelia burgdorferi*. Infect. Immun. 1992; 60 (12); 4995-5003
145. Kaneda K, Masuzawa T, Simon MM, Isogai E, Isogai H, Yasugami K, Suzuki T, Suzuki Y, Yanagihara Y. Infectivity and arthritis induction of *Borrelia japonica* on SCID mice and immune competent mice: possible role of galactosylceramide binding activity on initiation of infection. Microbiol. Immunol. 1998; 42 (3); 171-175
146. Weller M, Stevens A, Sommer N, Wietholter H. Are CSF or serum ganglioside antibodies related to peripheral nerve demyelination in neuroborreliosis, Guillain-Barre syndrome, or chronic inflammatory demyelinating polyradiculoneuropathy? Eur. Arch. Psychiatry Clin. Neurosci. 1992; 242 (2-3); 122-126
147. Weller M, Stevens A, Sommer N, Dichgans J, Kappler B, Wietholter H. Ganglioside antibodies: a lack of diagnostic specificity and clinical utility? J. Neurol. 1992; 239 (8); 455-459
148. Garcia-Monco JC, Wheeler CM, Benach JL, Furie RA, Lukehart SA, Stanek G, Steere AC. Reactivity of neuroborreliosis patients (Lyme disease) to cardiolipin and gangliosides. J. Neurol. Sci. 1993; 117 (1-2); 206-214
149. Garcia-Monco JC, Seidman RJ, Benach JL. Experimental immunization with *Borrelia burgdorferi* induces development of antibodies to gangliosides. Infect. Immun. 1995; 63 (10); 4130-4137
150. Burgdorfer W. Lyme-Borreliosis. MMW 1989; 131; 91-98
151. Hassler D, Simon MM, Kramer MD. Lyme-Borreliose. Neue Aspekte zum Verständnis von Pathophysiologie und Epidemiologie. Notabene Medici 1989; 718; 333-338
152. Wilske B, Prae-Mursic V, Schierz G. Lyme-Borreliose. Die gelben Hefte 1988; 27; 146-159
153. Ackermann R. Erythema migrans – Borreliose und Frühsommer-meningoenzephalitis. Dtsch. Ärzteblatt 1986; 83; 1765-1774
154. Weber K, Neubert U. Clinical features of early erythema migrans disease and related disorders. Zbl. Bakt. Hyg. 1986; (A263); 209-228
155. Weber K, Schierz G, Wilske B, Prae-Mursic V. Das Lymphozytom – eine Borreliose? Z. Hautkr. 1985; 60; 1585-1598

-
156. Albrecht S, Hofstadter S, Artsob H, Chaban O, From L. Lymphadenosis benigna cutis resulting from *Borrelia* infection (*Borrelia lymphozytoma*). J. Amer. Acad. Dermatol. 1991; 24; 621-625
 157. Horst H. Klinik der *Borrelia burgdorferi*-Infektionen – allgemeiner Überblick. Horst, H.(Hrsg): Einheimische Zeckenborreliose (Lyme – Krankheit) bei Mensch und Tier. Perimed Erlangen 1991; 54-58
 158. Logigian EL, Kaplan RF, Steere AC. Chronic neurologic manifestations of Lyme disease. N. Engl. J. Med. 1990; 323; 1438-1444
 159. Kristoferitsch W. Neuropathien bei Lyme-Borreliose. Springer Verlag Wien/New York 1989.
 160. Jonsson L, Stiernstedt G, Thomander L. Tick borne *Borrelia* infection in patients with Bell's palsy. Arch. Otolaryngol. Head Neck Surg. 1987; 113; 303-306
 161. McAlister HF, Klementowicz PT, Andrews C, Fisher JD, Feld M, Furman S. Lyme carditis: An important cause of atrioventricular heart block. Ann. Intern. Med. 1989; 110; 339-345
 162. Stanek G, Klein J, Bittner R, Glogar D. Isolation of *Borrelia burgdorferi* from the myocardium of a patient with longstanding cardiomyopathy. N. Engl. J. Med. 1990; 322; 249-252
 163. Jacobson DM, Marx JJ, Dlesk A. Frequency and clinical significance of Lyme seropositivity in patients with isolated optic neuritis. Neurol. 1991; 41; 706-711
 164. Duray P, Steere AC. The spectrum of organ and systems pathology in human Lyme disease. Zbl. Bakt. Hyg. 1986; A263; 169-178
 165. Bianchi G, Rovetta G, Monteforte P, Fumarola D, Trevisan G, Crovto F, Cimmino MA. Articular involvement in European patients with Lyme disease. A report of 32 Italian patients. Brit. J. Rheumatol. 1990; 29, 178-180
 166. Cruz MH, Hansen K, Faarvang KL, Link H. Intrasyovial production of oligoclonal specific antibodies in Lyme arthritis. N. Engl. J. Med. 1990; 322; 1889-1890
 167. Kramer MD, Moter SE, Hofmann H, Schaible UE, Simon MM, Wallich R. Symptomatik und Diagnostik der Lyme-Borreliose. Dtsch. Med. Wschr. 1993; 118; 423-427
 168. Asbrink E, Brehmer-Anderson E, Hovmark A. Acrodermatitis chronica atrophicans – A spirochetosis. Clinical and histopathological picture based on 32 patients; course and

- relationship to erythema chronicum migrans Afzelius. Amer. J. dermatopath. 1986; 8; 209-219
169. Aberer E, Stanek G, Ertl M, Neumann R. Evidence of spirochetal origin of circumscribed scleroderma (morphea). Acta. Derm. Venereol. (Stockh.) 1987; 67; 225-231
170. Aberer E, Stanek G. Histological evidence for spirochetal origin of morphea and lichen sclerosus et atrophicans. Amer. J. Dermatopath. 1987; 9; 374-379
171. Tuffanelli D. Do some patients with morphea and lichen sclerosus et atrophicans have a *Borrelia* infection? Amer. J. Dermatopath. 1987; 9; 371-373
172. Dickman S. Possible cause found for Lyme arthritis. Science 1998; 281 (5377); 631-632
173. Fendler C, Wu P, Eggens U, Laitko S, Sorenson H, Disler A, Braun J, Sieper J. Longitudinal investigation of bacterium-specific synovial lymphocyte proliferation in reactive arthritis and lyme arthritis. Br. J. Rheumatol. 1998; 37 (7); 784-788
174. Akin E, McHugh GL, Flavell RA, Fikrig E, Steere AC. The immunoglobulin (IgG) antibody response to OspA and OspB correlates with severe and prolonged Lyme arthritis and the IgG response to P35 correlates with mild and brief arthritis. Infect. Immun. 1999; 67 (1); 173-181
175. Linssen WH, Gabreels FJ, Wevers RA. Infective acute tranverse myelopathy. Report of two cases. Neuropediatrics 1991; 22 (2); 107-109
176. Garcia-Monco JC, Benach JL. Mechanisms of injury in Lyme neuroborreliosis. Semin. Neurol. 1997; 17 (1); 57-62
177. Strobino BA, Williams CL, Abid S, Chalson R, Spierling P. Lyme disease and pregnancy outcome: a prospective study of two thousand prenatal patients. Am. J. Obstet. Gynecol. 1993; 169; 367-374
178. Silver RM, Yang L, Daynes RA, Branch DW, Salafia CM, Weis JJ. Fetal outcome in murine Lyme disease. Infect. Immun. 1995; 63 (1); 66-72
179. Liebisch G, Sohns B, Bautsch W. Detection and typing of *Borrelia burgdorferi* sensu lato in Ixodes ricinus ticks attached to human skin by PCR. J. Clin. Microbiol. 1998; 36 (11), 3355-3358
180. Eiffert H, Karsten A, Thompsen R, Christen HJ. Characterization of *Borrelia burgdorferi* strains in Lyme arthritis. Scan. J. Infect. Dis. 1998; 30 (0); 265-268
181. Vasiliu V, Herzer P, Rossler DE, Lehnert G, Wilske B. Heterogeneity of *Borrelia burgdorferi* sensu lato demonstrated by an ospA-type-specific PCR in synovial fluid

- from patients with Lyme arthritis. *Med. Microbiol. Immunol. (Berlin)* 1998; 187 (2); 97-102
182. von Orelli S, Schnarwyler B, Maurer R, Hirsch H, von Castelberg B. Vulvares pseudolymphom: Nachweis der Infektion mit *Borrelia burgdorferi* mittels Polymerasekettenreaktion. *Gynäkol. Geburtshilfliche Rundsch.* 1998; 38 (3); 143-145
183. Liveris D, Varde S, Iyer R, Koenig S, Bittker S, Cooper D, McKenna D, Nowakowski J, Nadelman RB, Wormser GP, Schwartz I. Genetic Diversity of *Borrelia burgdorferi* in Lyme disease patients as determined by culture versus direct PCR with clinical specimens. *J. Clin. Microbiol.* 1999; 37 (3); 565-569
184. Brettschneider S, Bruckbauer H, Klugbauer N, Hofmann H. Diagnostic value of PCR for detection of *Borrelia burgdorferi* in skin biopsy and urine sample from patients with skin borreliosis. *J. Clin. Microbiol.* 1998; 36 (9); 2658-2665
185. Lebech AM, Hansen K, Rutledge BJ, Kolbert CP, Rys PN, Persing DH. Diagnostic detection and direct genotyping of *Borrelia burgdorferi* regular by Polymerase Chain Reaction in cerebrospinal fluid in Lyme neuroborreliosis. *Mol Diagn.* 1998; 3 (3); 131-141.
186. Pahl A, Kuhlbrandt U, Brunne K, Rollinghoff M, Gessner A. Quantitative detection of *Borrelia burgdorferi* by real-time PCR. *J. Clin Microbiol.* 1999; 37 (6); 1958-1963.
187. Barbour AG, Tessier SL, Todd WJ. Lyme disease spirochetes and ixodid tick spirochetes share a common surface antigenic determinant defined by a monoclonal antibody. *Infect. Immun.* 1983; 41; 795-804
188. Barbour AG, Tessier SL, Hayes SF. Variation in a major surface protein of Lyme disease spirochetes. *Infect. Immun.* 1984; 45; 94-100
189. Fikrig E, Barthold SW, Marcantonio N, Deponte K, Kantor FS, Flavell RA. Roles of OspA, OspB and flagellin in protective immunity to Lyme borreliosis in laboratory mice. *Infect. Immun.* 1992; 60; 657-661
190. Langermann S, Palaszinski S, Sadziene A, Stover CK, Koenig S. Systemic and mucosal immunity induced by BCG vector expressing outer surface protein A of *Borrelia burgdorferi*. *Nature* 1995; 372; 552-555
191. Shanafelt M, Anzola J, Soderberg C, Yssel H, Turck CW, Peltz G. Epitopes on the outer surface protein A of *Borrelia burgdorferi* recognized by antibodies and T cells of patients with Lyme disease. *Eur. J. Immunol.* 1992; 148; 218-224

-
192. Honarvar N, Schaible UE, Wallich R, Simon MM. A 14.000 MW lipoprotein and a glycolipid-structure of *Borrelia burgdorferi* induce proliferation and immunoglobulin production in mouse B cells at high frequencies. *Immunology* 1994; 82; 389-396
 193. Weis JJ, Ma Y, Erdile LF. Biological activities of native and recombinant *Borrelia burgdorferi* outer surface protein A: dependence on lipid modification. *Infect. Immun.* 1994; 62; 4632-4636
 194. Comstock LE, Thomas DD. Penetration of endothel cell monolayers by *Borrelia burgdorferi*. *Infect. Immun.* 1989; 57; 1626-1628
 195. Sadziene A, Barbour AG, Rosa PA, Thomas DD. An OspB mutant of *Borrelia burgdorferi* has reduced invasiveness in vitro and reduced infectivity in vivo. *Infect. Immun.* 1993; 61; 3590-3596
 196. Sadziene A, Thomas DD, Barbour AG. *Borrelia burgdorferi* mutant lacking Osp: biological and immunological characterization. *Infect. Immun.* 1995; 63; 1573-1580
 197. Thomas DD, Comstock LE. Interaction of Lyme disease spirochetes with cultured eucaryotic cells. *Infect. Immun.* 1989; 57; 1324-1326
 198. Barbour AG, Heiland RA, Howe TR. Heterogeneity of major proteins in Lyme disease borreliae: a molecular analysis of north american and european isolates. *J. Infect. Dis.* 1985; 152; 478-484
 199. Peter O, Bretz AG. Polymorphism of outer surface proteins of *Borrelia burgdorferi* as a tool for classification. *Zbl. Bakt.* 1992; 277; 28-33
 200. Pachner AR, Itano A, Ricalton N. An outer surface protein is a major T cell antigen in experimental Lyme disease in CB6F1 mice. *Ann. NY. Acad. Sci.* 1991; 636; 393-395
 201. Wilske B, Prae-Mursic, Schierz G, Busche KV. Immunochemical and immunological analysis of european *Borrelia burgdorferi* strains. *Zbl. Bakt. Hyg* 1986; A263; 92-102
 202. Luft BJ, Jiang W, Munoz P, Dattwyler RJ, Gorevic PD. Biochemical and immunological characterization of the surface proteins of *Borrelia burgdorferi*. *Infect. Immun.* 1989; 57; 3637-3645
 203. Bunikis J, Olsen B, Westman G, Bergsröm S. Variable serum immunoglobulin responses against different *Borrelia burgdorferi* sensu lato species in a population at risk for and patients with Lyme disease. *J. clin. Microbiol.* 1995; 33; 1473-1478
 204. Szczepanski A, Benach JL. Lyme-Borreliosis: host responses to *Borrelia burgdorferi*. *Micrbiol. Rev.* 1991; 55; 21-34

-
205. Sadziene A, Thompson PA, Barbour AG. In vitro inhibition of *Borrelia burgdorferi* growth by antibodies. J. Infect. Dis. 1993; 167; 165-172
 206. Probert WS, Allsup KM, LeFebvre RB. Identification and characterization of a surface-exposed, 66-kilodalton protein from *Borrelia burgdorferi*. Infect. Immun. 1995; 63; 1933-1939
 207. Rasiah C, Schlitz E, Reichert J, Vogt A. Purification and characterization of a tryptic peptide of *Borrelia burgdorferi* flagellin, which reduces cross-reactivity in immunoblots and ELISA. J. Gen. Microbiol. 1992; 138; 147-154
 208. Schneider T, Lange R, Röspeck W, Weigelt W, Kölmel HW. Prognostic B-cell epitopes on the flagellar protein of *Borrelia burgdorferi*. Infect. Immun. 1992; 60; 316-319
 209. Benach JL, Coleman JL, Garcia-Monco JC, Deponte PC. Biological activity of *Borrelia burgdorferi* antigens. Ann. NY Acad. Sci. 1988; 539; 115-125
 210. Harris A, Rosenberg AA, Riedel LM. A microflocculation test for syphilis using cardiolipin antigen. J. Vener. Dis. Inf. 1946;
 211. Pangborn MC, Almeida JO, Maltaner F, Siverstein AM, Thompson WR. Cardiolipin antigens. WHO, Monograph Series, Geneva 1955.
 212. Portnoy J. Modification of the rapid plasma (RPR) card test for syphilis for use in large scale testing. Am. J. Clin. Pathol. 1963; 40; 473-479
 213. U.S. Department of Health, Education and Welfare, National Communicable Center. Manual of Test of Syphilis. Venereal Disease Program 1969
 214. Coffey EM, Bradford L. Serodiagnosis of Syphilis. Manual Clinical Immunology 1980; 74; 530-541
 215. Steere AC. Lyme disease – a growing threat to urban populations. Proc. Natl. Acad. Sci. USA 1994; 91; 2378-2383
 216. Kramer MD, Wallich R, Simon MM. Mechanismen der Pathogenese und Protektion bei der Lyme-Borreliose. Süss J (Hrsg.): Durch Zecken übertragbare Erkrankungen – FSME und Lyme-Borreliose. 3. Potsdamer Symposium 1995. Weller Verlag
 217. Alonso-Llamazares J, Persing DH, Anda P, Gibson LE, Rutledge BJ, Iglesias L. No evidence for *Borrelia burgdorferi* infection in lesions of morphea and lichen sclerosus et atrophicus in Spain. A prospective study and literature review. Acta Derm. Venereol. 1977; 77 (4); 299-304

-
218. Matthews HM, Yang TK, Jenkin HM. Alk-1-enyl ether phospholipids (plasmalogens) and glycolipids of *Treponema hyodysenteriae*. *Biochemica et Biophysica Acta* 1980; 618; 273-281
 219. Morrison DC, Jacobs DM. Lipopolysaccharides and endotoxin. In: Sirois P, Rola-Pleszczynski M. *Immunopharmacology*. Amsterdam: Elsevier Biomedical Press 1982: 169-197
 220. Mackworth-Young CG, Harris EN, Steere AC et al. Anticardiolipn antibodies in Lyme Disease. *Arthritis Rheum.* 1989; 31; 1052-1056
 221. Pedersen NS, Orum O, Mouritsen S. Enzyme-linked immunoabsorbent assay for detection of antibodies to the Veneral Disease Research Laboratory (VDRL) antigen in syphilis. *J. clin. Microbiol.* 1987; 25; 1711-1716
 222. Garcia-Monco JC, Fernandez Villar B, Benach JL. Adherence of the Lyme disease spirochete to glial cells and the cells of glial origin. *J. Infect. Dis.* 1989; 160; 497-506
 223. Isaacs, RD. *Borrelia burgdorferi* bind to epithelial cell proteoglycans. *J. Clin. Invest.* 1994; 93; 809-819
 224. Stults CLM, Sweeley CC, Macher BA. Glycosphingolipids: structure, biological source, and properties. *Methods Enzymol.* 1989; 179; 167-214
 225. Garcia-Monco JC, Fernandez Villar B, Szczepanski A, Benach JL. Cytotoxicity of *Borrelia burgdorferi* for cultured rat glial cells. *J. Infect. Dis.* 1991; 163; 1362-1366
 226. Wirguin I, Suturkova-Milosevic L, Della-Latta P, Fisher T, Brown RH, Latov N. Monoclonal IgM antibodies to GM1 and asialo-GM1 in chronic neuropathies cross-react with *Campylobacter jejuni* lipopolysaccharides. *Ann. Neurol.* 1994; 35; 698-703
 227. Mishu B, Blaser MJ. Role of infection due to *Campylobacter jejuni* in the initiation of Guillain-Barre-Syndrome. *Clin. Infect. Dis.* 1993; 17; 101-108
 228. Radolf JD, Arndt LL, Akins DR, Curetty LL, Levi ME, Shen Y, Davis LS, Norgard MV. *Treponema pallidum* and *Borrelia burgdorferi* lipoproteins and synthetic lipopeptides activate monocyte/macrophages. *J. Immunol.* 1995; 154; 2866-2877
 229. Platt FM, Butters TD. New therapeutic prospects for the glycosphingolipid lysosomal storage diseases. *Biochem Pharmacol.* 1998 Review; 15;56 (4);421-30.
 230. Hirano T, Hashimoto H, Shiokawa Y, Iwamori M, Nagai Y, Kasai M, Ochiai Y, Okumura K. Antiglycolipid antibody detected in the sera from systemic lupus erythematosus patients. *J. Clin. Invest.* 1980; 66; 1437; 1440

-
231. Gleeson PA. Glycoconjugates in autoimmunity. *Biochim. Biophys. Acta* 1994; 1197; 237-255
232. Shaw N. Bacterial glycolipids and glycophospholipids. *Adv. Microbio. Physiol.* 1975; 12; 141-167
233. Fischer W. Newly discovered lipids from Streptococci. *Lipids* (R. Paoletti, G. Porcellatti, G. Jacini, Ed.) 1976; Raven Press, NY; 255-266
234. Marchase RB. Biochemical investigations of retinotectal adhesive specificity. *J. Cell. Biol.* 1977; 75; 237-257
235. Karlsson KA. Animal glycosphingolipids as membrane attachment sites for bacteria. *Ann. Rev. Biochem.* 1989; 58, 309-350
236. Adams GA, Singh PP. Structural features of Lipid A preparations isolated from *E. coli* and *S. flexneri*. *Biochim. Biophys. Acta* 1970; 202; 553-555
237. Rietschel ET, Kirikae T, Schade FU, Brade H et al. Bacterial endotoxin: molecular relationships of structure to activity and function. *FASEB J.* 1994; 8; 217-225
238. Rietschel ET, Brade H, Holst O, Schumann RR et al. Bacterial endotoxin: chemical constitution, biological recognition, host response, and immunological detoxification. *Curr. Topics in Microbiol. Immunol* (E.T. Rietschel, H. Wagner, Ed.) 1996; 40-81; Springer-Verlag, Berlin
239. Kiarash A, Boyd B, Lingwood CA. Glycosphingolipid receptor function is modified by fatty acid content. *J. Biol. Chem.* 1994; 1269; 11138-11146
240. Hamilton KS, Briere K, Jarrell HC, Grant CWM. Acyl chain length effects related to glycosphingolipid crypticity in phospholipid membrane: probed by 2H-NMR. *Biochim. Biophys. Acta* 1994; 1190; 367-375
241. Satz N. Klinik der Lyme-Borreliose. Verlag Hans Huber 1993; 12
242. Hovind-Hougen K, Asbrink E, Stiernstedt G, Steere AC, Hovmark A. Ultrastructural differences among spirochetes isolated from patients with Lyme disease and related disorders, and from *Ixodes ricinus*. *Zbl. Bakt. Hyg.* 1986; A263; 103-111
243. Gray JS, Kahl O, Robertson JN, Daniel M, Estrada-Pena A, Gettinby G, Jaenson TG, Jensen P, Jongejan F, Korenberg E, Kurtenbach K, Zeman P. Lyme borreliosis habitat assessment. *Zbl. Bakteriologie*. 1998; 287 (3):211-28.
244. Gylfe, Olsen B, Strasevicius D, Marti Ras N, Weihe P, Noppa L, Ostberg Y, Baranton G, Bergstrom S. Isolation of Lyme disease *Borrelia* from puffins (*Fratercula arctica*)

-
- and seabird ticks (*Ixodes uriae*) on the Faeroe Islands. J. Clin. Microbiol. 1999; 37 (4): 890-896
245. Lehninger A. Der supramolekulare Bau von Membranen. Aus: Prinzipien der Biochemie (Lehninger, Nelson, Cox; ED.), 2. Auflage 1994; Seite 315; Spektrum Akademischer Verlag, Heidelberg