

## 7 References:

1. McKusick VA. Spastic paraplegia 3A. Online mendelian Inheritance in Man, OMIM (TM). McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, MD)2000. World Wide Web URL: [Http://www.ncbi.nlm.nih.gov/omim/](http://www.ncbi.nlm.nih.gov/omim/)
2. Fink JK, Heiman-Patterson T, the Hereditary Spastic Paraplegia Working Group. Hereditary spastic paraplegia: advances in genetic research. *Neurology* 1996; **335**: 1169-1175.
3. Baraitser M. Spastic paraplegia/HSP. In: Motulsky AG, Bobrow M, Harper PS, Scriver C, eds. *The genetics of neurological disorders*. 2<sup>nd</sup> ed. New York: Oxford University Press, 1990:275-290.
4. Harding AE. Hereditary spastic paraplegias. *Semin Neurol* 1993; **13**: 333-336.
5. Werderlin L. Hereditary ataxias. Occurrence and clinical features. *Acta Neurol Scand* 1986; **73** (suppl. 106): 124.
6. Skre H. Hereditary spastic paraplegia in western Norway. *Clin Genet* 1974; **6**: 165-183.
7. Polo JM, Calleja J, Combarros O. Hereditary ataxias and paraplegias in Cantabria, Spain. An epidemiological and clinical study. *Brain* 1991; **114**: 855-866.
8. Reid E. Pure hereditary spastic paraplegia. *J Med Genet* 1997; **34**: 499-503.
9. Harding AE. Hereditary "pure" spastic paraplegia: a clinical and genetic study of 22 families. *J Neurol Neurosurg Psychiatry* 1981; **44**: 871-883.
10. Keppen LD, Leppert MF, O'Connell PO. Etiological heterogeneity in X-linked spastic paraplegia. *Am J Hum Genet* 1987; **41**: 933-943.
11. Strümpell A. Beiträge zur Pathologie des Rückenmarks. *Arch Psychiatr Nervenkr* 1880; **10**: 676-717.

12. Dürr A, Brice A, Serdaru M, Rancurel G, Derouesne C, Lyon-Caen O, Agid Y, Fontaine B. The phenotype of "pure" autosomal dominant spastic paraplegia. *Neurology* 1994; **44**: 1274-1277.
13. Behan W, Maia M. Strumpell's familial spastic paraplegia genetics and neuropathology. *J Neurol Neurosurg Psychiatry* 1974; **37**: 8-20.
14. Kenwrick S, Ionasescu V, Ionasescu G, Searby C, King A, Dubowitz M, Davies KE. Linkage studies of X-linked recessive spastic paraplegia using DNA probes. *Hum Genet* 1986; **73**: 264-266.
15. Winter RM, Davies KE, Bell MV, Huson SM, Patterson MN. MASA syndrome: further clinical delineation and chromosomal localisation. *Hum Genet* 1989; **82**: 367-370.
16. Schrandt-Stumpel C, Legius E, Fryns JP, Cassiman JJ. MASA syndrome: new clinical features and linkage analysis using DNA probes. *J Med Genet* 1990; **27**: 688-692.
17. Jouet M, Rosenthal A, Armstrong G, MacFarlane J, Stevenson R, Paterson J, Metzenberg A, Ionasescu V, Temple K, Kenwrick S. X-linked spastic paraplegia (SPG1), MASA syndrome and X-linked hydrocephalus result from mutations in the L1 gene. *Nature Genet* 1994; **7**: 402-407.
18. Bonneau D, Rozet JM, Bulteau C, Berthier M, Mettey R, Gil R, Munnich A, Le Merrer M. X linked spastic paraplegia (SPG2): clinical heterogeneity at a single gene locus. *J Med Genet* 1993; **30**: 381-384.
19. Saugier-Verber P, Munnich A, Bonneau D, Rozet JM, Le Merrer M, Gil R, Boespflug-Tanguy O. X-linked spastic paraplegia and Pelizaeus-Merzbacher disease are allelic disorders at the proteolipid protein locus. *Nature Genet* 1994; **6**: 257-262.
20. Kobayashi H, Hoffman EP, Marks HG. The rumpshaker mutation in spastic paraplegia. (Letter) *Nature Genet* 1994; **7**: 351-352.

- 
21. Schneider A, Montague P, Griffiths I, Fanarraga M, Kennedy P, Brophy P, Nave KA. A mutation of the proteolipid protein gene uncoupling hypomyelination from glial cell death. *Nature* 1999; **358**: 758-761
  22. Steinmuller R, Lantigua-Cruz A, Garcia-Garcia R, Kostrzewa M, Steinberger D, Müller U. Evidence of a third locus in X-linked recessive spastic paraplegia. *Hum Genet* 1997; **100**: 287-289.
  23. Hentati A, Pericak-Vance MA, Hung WY, Belal S, Laing N, Boustany RM, Hentati F, Ben Hamida M, Siddique T. Linkage of 'pure' autosomal recessive familial spastic paraplegia to chromosome 8 markers and evidence of genetic locus heterogeneity. *Hum Mol Genet* 1994; **3**: 1263-1267.
  24. De Michele G, De Fusco M, Cavalcanti F, Filla A, Marconi R, Volpe G, Monticelli A, Ballabio A, Casari G, Coccozza S. A new locus for autosomal recessive hereditary spastic paraplegia maps to chromosome 16q24.3. *Am J Hum Genet* 1998; **63**: 135-139.
  25. Casari G, De Fusco M, Ciarmatori S, Zeviani M, Mora M, Fernandez P, De Michele G, Filla A, Coccozza S, Marconi R, Dürr A, Fontaine B, Ballabio A. Spastic paraplegia and OXPHOS impairment caused by mutations in paraplegin, a nuclear-encoded mitochondrial metalloprotease. *Cell* 1998; **93**: 973-983.
  26. Confalonieri F, Duguet M. A 200-amino acid ATPase module in search of a basic function. *Bioessays* 1995; **17**: 639-650.
  27. Patel S, Latterich M. The AAA team: related ATPases with diverse functions. *Trends Cell Biol* 1998; **8**: 65-71.
  28. Neuwald AF, Aravind L, Spouge JL, Koonin EV. AAA<sup>+</sup>: a class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes. *Genome Res* 1999; **9**: 27-43.

- 
29. Martinez Murillo F, Kobayashi H, Pegoraro E, Galluzzi G, Creel G, Mariani C, Farina E, Ricci E, Alfonso G, Pauli RM, Hoffman EP. Genetic localization of a new locus for recessive familial spastic paraparesis to 15q13-15. *Neurology* 1999; **53**: 50-56.
30. Vazza G, Zortea M, Boaretto F, Micaglio GF, Sartori V, Mostacciuolo, ML. A new locus for autosomal recessive spastic paraplegia associated with mental retardation and distal motor neuropathy, SPG14, maps to chromosome 3q27-q28. *Am J Hum Genet* 2000; **67**: 504-509.
31. Hazan J, Lamy C, Melki J, Munnich A, de Recondo J, Weissenbach J. Autosomal dominant familial spastic paraplegia is genetically heterogeneous and one locus maps to chromosome 14q. *Nature Genet* 1993; **5**: 163-167.
32. Hentati A, Pericak-Vance MA, Lennon F, Wasserman B, Hentati F, Juneja T, Angrist MH, Hung WY, Boustany RM, Bohlega S, *et al.* Linkage of a locus for autosomal dominant familial spastic paraplegia to chromosome 2p marker. *Hum Mol Genet* 1994; **3**: 1867-1871.
33. Gispert S, Santos N, Damen R, Orozco G, Ha-Hao D, Ratzlaff T, Aguiar J, Torrens I, Heredero L, Brice A, *et al.* Autosomal dominant familial spastic paraplegia: reduction of the FSP1 candidate region chromosome 14q to 7cM and locus heterogeneity. *Am J Hum Genet* 1995; **56**: 183-187.
34. Hazan J, Fontaine B, Bruyn RPM, Lamy C, van Deutekom JCT, Rime CS, Dürr A, Melki J, Lyon-Caen O, Agid Y, Munnich A, Padberg GW, de Recondo J, Frants RR, Brice A, Weissenbach J. Linkage of a new locus for autosomal dominant familial spastic paraplegia to chromosome 2p. *Hum Mol Genet* 1994; **3**: 1569-1573.
35. Scott WK, Gaskell PC, Lennon F, Wolpert CM, Menold MM, Aylsworth AS, Warner C, Farrell CD, Boustany RMN, Albright SG, Boyd E, Kingston HM, Cumming WJK, Vance JM, Pericak-Vance MA. Locus heterogeneity, anticipation and reduction of the chromosome 2p minimal candidate region in autosomal dominant familial spastic paraplegia. *Neurogenetics* 1997; **1**: 95-102.

- 
36. Hazan J, Fonknechten N, Mavel D, Paternotte C, Samson D, Artiguenave F, Davoine CS, Cruaud C, Dürr A, Wincker P, Brottier P, Cattolico L, Barbe V, Burgunder JM, Prud'homme JF, Brice A, Fontaine B, Heilig R, Weissenbach J. Spastin, a new AAA protein, is altered in the most frequent form of autosomal dominant spastic paraplegia. *Nature Genet* 1999; **23**: 296-303.
37. Hazan, J, Davoine C-S, Mavel D, Fonknechten N, Paternotte C, Fizames C, Cruaud C, Damson D, Muselet D, Vega-Czarny N, Brice A, Gyapay G, Heilig R, Fontaine B, Weissenbach J. A Fine intergarted map of the SPG4 locus excludes an expanded CAG repeat in chromosome 2p-linked autosomal dominant spastic paraplegia. *Genomics* 1999; **60**: 309-319.
38. Fink JK, Wu CB, Jones SM, Sharp GB, Lange BM, Lesicki A, Reinglass T, Varvil T, Otterud B, Leppert M. Autosomal dominant familial spastic paraplegia: tight linkage to chromosome 15q. *Am J Hum Genet* 1995 ; **56**: 188-192.
39. Hedera, P, Rainier S, Alvarado D, Zhao X, Williamson J, Otterud B, Leppert M, Fink JK. Novel locus for autosomal dominant hereditary spastic paraplegia, on chromosome 8q. *Am J Hum Genet* 1999 ; **64**: 563-569.
40. Seri M, Cusano R, Forabosco P, Cinti R, Caroli F, Picco P, Bini R, Morra VB, De Michele G, Lerone M, Silengo M, Pela I, Borrone C, Romeo G, Devoto M. Genetic mapping to 10q23.3-q24.2, in a large Italian pedigree, of a new syndrome showing bilateral cataracts, gastroesophageal reflux, and spastic paraparesis with amyotrophy. *Am J Hum Genet* 1999; **64**: 586-593.
41. Reid E, Dearlove AM, Rhodes M, Rubinsztein DC. A new locus for autosomal dominant 'pure' hereditary spastic paraplegia mapping to chromosome 12q13, and evidence for further genetic heterogeneity. *Am J Hum Genet* 1999; **65**: 757-763.

- 
42. Reid E, Dearlove AM, Osborn O, Rogers MT, Rubinsztein DC. A locus for autosomal dominant 'pure' hereditary spastic paraplegia maps to chromosome 19q13. *Am J Hum Genet* 2000; **66**: 728-732.
  43. Reid E, Grayson C, Rogers MT, Rubinsztein DC. Locus-phenotype correlations in autosomal dominant pure hereditary spastic paraplegia: a clinical and molecular genetic study of 28 United kingdom families. *Brain* 1999; **122**: 1741-1755.
  44. Fontaine B, Davoine CS, Dürr A, Paternotte C, Feki I, Weissenbach J, Hazan J, Brice A. A new locus for autosomal dominant pure spastic paraplegia, on chromosome 2q24-q34. *Am J Hum Genet* 2000; **66**: 702-707.
  45. Dracopoli NC, Haines JL, Korf BR, Moir DT, Morton CC, Seidman CE, Seidman JG, Smith DR. *Current Protocol in Human Genetics*. John Wiley & Sons, Inc. 1994.
  46. Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K. *Current Protocol in Molecular Biology*. John Wiley & Sons, Inc. 1993.
  47. Dubé MP, Rouleau, GA, Kibar Z, Farlow MR, Ebers G, Harper P, Kolodny EH, Baumbach L, Figlewicz DA. Autosomal dominant hereditary spastic paraplegia: Linkage analysis of a heterogeneous trait. *Am J Hum Genet* 1995; **57**: A190.
  48. Hudson TJ, Stein LD, Gerety SS, Ma J, Castle AB, Silva J, Slonim DK, Baptista R, Kruglyak L, Xu SH, *et al.* An STS-Based Map of the Human Genome. *Science* 1995; **270**: 1945-1954.
  49. De Jonghe P, Krols L, Michalik A, Hazan J, Smeyers G, Löfgren A, Weissenbach J, Martin JJ, Van Broeckhoven C. Pure Familial Spastic Paraplegia: Clinical and Genetic Analysis of Nine Belgian Pedigrees. *Eur J Hum Genet* 1996; **4**: 260-266.
  50. Lau EL, Kostrzewa M, Müller U. Transcript map of the chromosome 2 linked autosomal dominant spastic paraplegia (SPG4) critical region and identification of a highly informative STRP. *Neurogenetics* 1998; **2**: 75-76.

- 
51. Nielsen JE, Koefoed P, Abell K, Hasholt L, Eiber H, Fenger K, Niebuhr E, Sørensen SA. CAG repeat expansion in autosomal dominant pure spastic paraplegia linked to chromosome 2p21-p24. *Hum Mol Genet* 1997; **6**: 1811-1816.
52. Schalling M, Hudson TJ, Buetow KH, Housman DE. Direct detection of novel expanded trinucleotide repeats in the human genome. *Nature Genet* 1993; **4**: 135-139.
53. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997; **25**: 3389-3402.
54. Ichida K *et al.* Cloning of the cDNA encoding human xanthine dehydrogenase (oxidase): structural analysis of the protein and chromosomal location of the gene. *Gene* 1993; **133**: 279-284.
55. Kanzaki, T. *et al.* TGF- $\beta$  1 binding protein: a component of the large latent complex of TGF- $\beta$  1 with multiple repeat sequences. *Cell* 1990; **61**: 1051-1061.
56. Pohl U, Smith JS, Tachibana I, Ueki K, Lee HK, Ramaswamy S, Wu Q, Mohrenweiser HW, Jenkins RB, Louis DN. EHD2, EHD3, and eHD4 encode novel members of highly conserved family of EH domain-containing proteins. *Genomics* 2000; **63**: 255-262.
57. Collins FS. Positional cloning moves from predential to traditional. *Nature Genet.* 1995; **9**: 347-350.
58. Morgan TH. Random segregation versus coupling in Mendelian inheritance. *Science* 1911; **34**: 384.
59. Bell J, Haldane JBS. The linkage between the genes for color-blindness and haemophilia in men. *Proc R Soc Ser [Biol]* 1937; **123**: 119-150.

- 
60. Haldane JBS. The combination of linkage values and the calculation of distances between the loci of linked factors. *J Genet* 1919; **8**: 299-309.
61. Terwilliger JD, Ott J. *Handbook of human genetic linkage*. 1994. Johns Hopkins University Press, Baltimore.
62. Ott J. Analysis of human genetic linkage, 3<sup>rd</sup> ed. 1999. Johns Hopkins University Press, Baltimore.
63. Koenig M, Hoffman EP, Bertelsom CJ, Monaco AP, Feener C, Kunkel LM. Complete cloning of the Duchenne muscular dystrophy (DMD) cDNA and preliminary genomic organization of the DMD gene in normal and affected individuals. *Cell* 1987; **50**: 509-517.
64. Lee WH, Bookstein R, Hong F, Young LJ, Shew JY, Lee EY. Human retinoblastoma susceptibility gene: Cloning, identification, and sequence. *Science* 1987; **235**: 1394-1399.
65. Walter MA, Spillett DJ, Thomas P, Weissenbach J, Goodfellow PN. A method for constructing radiation hybrid maps of whole genomes. *Nature Genet* 1994; **7**: 22-28.
66. Krawczak M, Schmidtke J. *DNA Fingerprinting*, 2<sup>nd</sup> ed. BIOS Scientific Publishers, Oxford.
67. Sternberg N. Bacteriophage P1 cloning system for the isolation, amplification, and recovery of DNA fragments as large as 100 kilobase pairs. *Proc Natl Acad Sci USA* 1990; **87**: 103-107.
68. Dilella, AG, Woo SLC. Cloning large segments of genomic DNA using cosmid vecotrs. *Methods Enzymol*. 1987; **152**: 199-212.
69. Shizuya H, Birren B, Kim UJ, Mancino V, Slepak T, Tachiiri Y, Simon M. Cloning and stable maintenance of 300-kilobase-pair fragments of human DNA in *Escherichia coli* using an F-factor-based vector. *Proc Natl Acad Sci USA* 1992; **89**: 8794-8797.



- 
70. Ioannou Pa, Amemiya CT, Garnes J, Kroisel PM, Shizuya H, Chen C, Batzer MA, de Jong PJ. P1-derived vector for the propagation of large human DNA fragments. *Nature Genet* 1994; **6**: 84-89.
71. Burke DT, Carle GF, Olson MV. Cloning of large segments of exogenous DNA into yeast by means of artificial chromosome vectors. *Science* 1987; **236**: 806-812.
72. Cohen D, Chumakov I, Weissbach J. A first-generation map of the human genome. *Nature* 1993; **366**: 698-701.
73. Strachan T, Read AP. *Human Molecular Genetics*. BIOS Scientific Publishers, 1999.
74. Raskind WH, Pericak-Vance MA, Lennon F, Wolff J, Lipe HP, Bird TD. Familial Spastic Paraparesis: Evaluation of locus heterogeneity, anticipation, and Haplotype mapping of the SPG locus on the short arm of chromosome 2. *Am J Med Genet* 1997; **74**: 26-36.
75. Kremer E, Pritchard M, Lynch M, Yu S, Holman K, Baker E, Warren ST, Schlessinger D, Sutherland GR, Richards RI. Mapping of DNA instability at the fragile X to a trinucleotide repeat sequence p(CCG)<sub>n</sub>. *Science* 1991; **252**: 1711-1714.
76. Knight SJL, Flanner AV, Hirst MC, Campbell L, Christodoulou Z, Phelps SR, Pointon J, Middleton-Price HR, Barnicoat A, Pembrey ME, Holland J, Oostra BA, Bobrow M, Davies KE. Trinucleotide repeat amplification and hypermethylation of a CpG island in FRAXE mental retardation. *Cell* 1993; **74**: 127-134.
77. Gacy AM, Goellner GM, Spiro C, Chen X, Gupta G, Bradbury EM, Dyer RB, Mikesell MJ, Yao JZ, Johnson AJ, Richter A, Melancon SB, McMurray CT. GAA instability in Friedreich's Ataxia shares a common, DNA-directed and intraallelic mechanism with other trinucleotide diseases. *Mol Cell* 1998; **1**: 583-93.
78. Brook JD, McCurrach ME, Harley HG, Buckler AJ, Church D, Aburatani H, Hunter K, Stanton VP, Thirion JP, Hudson T, Sohn R, Zelman B, Snell RG, Rundle SA, Crow S, Davies J, Shelbourne P, Buxton J, Jones C, Juvonen V, Johnson K, Harper PS, Shaw DJ,

- Housman DE. Molecular basis of myotonic dystrophy: Expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* 1992; **68**: 799-808.
79. The Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 1993; **72**: 971-983.
80. La Spada AR, Wilson EM, Lubahn DB, Harding AE, Fischbeck KH. Androgen receptor gene mutations in X-linked spinal and bulbar muscular atrophy. *Nature* 1991; **352**: 77-79.
81. Orr HT, Chung M, Banfi S, Kwiatkowski TJ Jr, Servadio A, Beaudet AL, McCall AE, Duvick LA, Ranum LPW, Zoghbi HY. Expansion of an unstable trinucleotide (CAG) repeat in spinocerebellar ataxia type 1. *Nature Genet* 1993; **4**: 221-226.
82. Pulst SM, Nechiporuk A, Nechiporuk T, Gisper S, Chen XN, Lopes-Cendes I, Pearlman S, Starkman S, Orozco-Diaz G, Lunkes A, De Jong P, Rouleau GA, Auburger G, Korenber JR, Figueroa C, Sahba S. Moderate expansion of a normally biallelic trinucleotide repeat in spinocerebellar ataxia type 2. *Nature Genet* 1996; **14**: 269-276.
83. Kawaguchi Y, Okamota Z, Zaniwaki M, Aizawa M, Inoue M, Katayama S, Kawakami H, Nakamura S, Nishimura M, Akiguchi I, Kimur J, Narumiya S, Kakizuka A. CAG expansions in a novel gene for Machado-Joseph disease at chromosome 14q32.1. *Nature Genet*; **8**: 221-228.
84. Zhuchenko O, Bailey J, Bonnen P, Ashizawa T, Stockton DW, Amos C, Dobyns WB, Subramony SH, Zoghbi HY, Lee CC. Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the  $\alpha_1A$ -voltage-dependent calcium channel. *Nature Genet* 1997; **15**: 62-69.

- 
85. David G, Abbas N, Stevanin G, Dürr A, Yvert G, Cancel G, Wever C, Imbert G, Saudou F, Antoniou E, Drabkin H, Gemmil R, Giunti P, Benomar A, Wood N, Buber M, Agid Y, Mandel JL, Brice A. Cloning of the SCA7 gene reveals a highly unstable CAG repeat expansion. *Nature Genet* 1997; **17**: 65-70.
86. Koide R, Ikeuchi T, Onodera O, Tanaka H, Igarashi S, Endo K, Takasashi H, Kondo R, Ishikawa A, Hayashi T, Saito M, Tmoda A, Miike T, Naito H, Ikuta F, Tsuji S. Unstable expansion of CAG repeat in hereditary dentatorubral-pallidoluysian atrophy (DRPLA). *Nature Genet* 1994; **6**: 9-13.
87. Benson KF, Horwitz M, Wolff J, Griend K, Thompson E, White S, Richards RI, Raskind WH, Bird TD. CAG repeat expansion in autosomal dominant familial spastic paraparesis: novel expansion in a subset of patients. *Hum Molec Genet* 1998; **7**: 1779-1786.
88. Breshel TS, McInnis MG, Margolis RL, Sirugo G, Corneliussen B, Simpson SG, McMahon FJ, Mackinnon DF, Xu JF, Pleasant N, Huo Y, Ashworth rG, Grundstrom C, Grundstrom T, Kidd KK, DePaulo JR, Ross CA. A novel, heritable expanding CTG repeat in an intron of the SEF2-1 gene on chromosome 18q21.1. *Hum Mol Genet* 1997; **6**: 1855-1863.
89. Nakamoto M, Takebayashi H, Kawaguchi Y, Narumiya Y, Taniwaki M, Nakamura Y, Ishikawa Y, Akiyuchi I, Kimura J, Kakizuka A. A CAG/CTG expansion in the normal population. *Nature Genet* 1997; **17**: 358-386.
90. Del-Favero J, Goossens D, De Jongh P, Benson K, Michalik A, Ban den Bossche D, Horwitz M, Van Broeckhoven C. Isolation of CAG/CTG repeats from within the chromosome 2p21-p24 locus for autosomal dominant spastic paraplegia (SPG4) by YAC fragmentation. *Hum Genet* 1999; **105**: 217-225.
91. Andersson S, Berman DM, Jenkins EP, Russel DW. Deletion of steroid 5' reductase 2 gene in male pseudohermaphroditism. *Nature* 1991; **354**: 159-161.

- 
92. Fonknechten N, Mavel D, Byrne P, Davvoine CS, Cruaud C, Boentsch D, Samson D, Coutinho P, Hutchinson M, McMonagle P, Burgunder JM, Tartaglione An, Heinzlef O, Feki I, Deufel T, Parfrey N, Brice A, Fontaine B, Prud'homme JF, Weissenbach J, Dürr A, Hazan J. Sepectrum of SPG4 mutations in autosomal dominant spastic paraplegia. *Hum Mol Genet* 2000; **9**: 637-644.
93. Bürger J, Fonknechten N, Hoeltzenbein M, Neumann L, Bratanoff E, Hazan J, Reis A. Hereditary spastic paraplegia cased by mutations in the SPG4 gene. *Eur J Hum Genet.* 2000; **8**: 771-776.
94. Lindsey JC, Lusher ME, McDermott CJ, White KD, Reid E, Rubinsztein DC, Bashir R, Hazan J, Shaw PJ, Bushby KMD. Mutation analysis of the spastin gene (*SPG4*) in patients with hereditary spastic paraparesis. *J Med Genet* 2000; **37**: 759-765.
95. Santorelli FM, Patrono C, Fortini D, Tessa A, Comanducci G, Bertini E, Pierallini A, Nmabile GA, Casali C. Intrafamilial variability in hereditary spastic paraplegia associated with an *SPG4* gene mutation. *Neurology* 2000; **55**: 702-705.
96. Hentati A, Deng HX, Zhai H, Chen W, Yang Y, Hung WY, Azim AC, Bohlega S, Tandan R, Warner C, Laing NG, Cambi F, Mitsumoto H, Roos RP, Boustany RM, Ben Hamida M, Hentati F, Siddique T. Novel mutations in spastin gene and absence of correlation with age at onset of symptoms. *Neurology* 2000; **55**: 1388-1390.
97. Lau EL, Müller U. 2000. *Human Mutation database*: accession number H971836.
98. Sheffield VC, Beck JS, Kwitek AE, Sandstrom DW, Stone EM. The sensitveity of single strand conformaltional polymorphism analysis for the dtection of single base substitutions. *Genocmics* 1993; **16**: 325-332.
99. Cotton RGH, Edkins E, Forrest S. 1998 *Mutation Detection: a Pratical Approach*. IRL Press. Oxford.

- 
100. Mashal RD, Koontz J, Sklar J. Detection of mutations by cleavage of DNA heteroduplexes with bacteriophage resolvases. *Nature Genet* 1995; **9**: 177-183.
101. van der Lijst R, Khan PM, Vaen H, van Leeuwen C, tops C, Roest PA et al. Rapid detection of translation-terminating mutations at the adenomatous polyposis coli (APC) gene by direct protein truncation test. *Genomics* 1994; **11**: 365-368.
102. Hacia JG. Resequencing and mutational analysis using oligonucleotide microarrays. *Nature Genet* 1999; **21** (suppl.): 42-47.
103. Chelly J, Kaplan JC, Maire P, Gautron S, Kahn A. Transcription of the dystrophin gene in human muscle and non-muscle tissue. *Nature* 1988; **333**: 858-860.
104. McMonagle P, Byrne PC, Fitzgerald B, Webb S, Parfrey NA, Hutchinson M. Phenotype of AD-HSP due to mutations in the SPAST gene. *Neurology* 2000; **55**: 1794-1800.
105. Hilt W, Wolf DH. Proteasomes of the yeast *S.cerevisiae*: genes, structure and functions. *Mol biol Reports* 1995; **21**: 3-10.
106. Liberzon A, Shpungin S, Bangion H, Yona E, Katcoff DJ. Association of yeast SAP1, a novel member of the ``AAA`` ATPase family of proteins, with the chromatin protein SIN1. *FEBS Lett* 1996; **388**: 5-10.
107. Andrews BJ, Mason SW. Gene expression and the cell cycle; a family affair. *Science* 1993; **261**: 1543-1544.
108. Andrews BJ, Donoviel MS. A heterodimeric transcriptional repressor becomes clear. *Science* 1995; **270**: 251-253.

- 
109. Ozelius LJ, Hewett JW, Page CE, Bressman SB, Krmer PL, Shalish C, de Deon D, Brin MF, Raymond D, Corey DP, Fahn S, Risch NJ, Buckler AJ, Gusella JF, Breakefield XO. The early-onset torsion dystonia gene (DYT1) encodes an ATP-binding protein. *Nature Genet.* 1997; **17**: 40-48.
110. Hardy J, Gwinn-Hardy K. Genetic classification of primary neurodegenerative disease. *Science* 1998; **282**: 1075-1079.
111. Heinzlef O, Paternotte C, Jean-François Prud'homme F M, Dien J, Madigand M, Puget J, Weissenbach J, Roullet E, Hazan J. Mapping of a complicated familial spastic paraplegia to locus SPG4 on chromosome 2p. *J Med Genet* 1998; **35**: 89-93.
112. Webb S, Coleman D, Byrne P, Parfrey N, Burke T, Hutchinson J, Hutchinson M. Autosomal dominant hereditary spastic paraparesis with cognitive loss linked to chromosome 2p. *Brain* 1998; **121**: 601-609.
113. White KD, Ince PG, Lusher M, Lindsey J, Cookson M, Bahir R, Shaw PJ, Bushby KMD. Clinical and pathologic findings in hereditary spastic paraparesis with spastin mutation. *Neurology* 2000; **55**: 89-94.
114. Byrne PC, Mc Monagle P, Webb S, Fitzgerald B, Parfrey NA, Hutchinson M. Age-related cognitive decline in hereditary spastic paraparesis linked to chromosome 2p. *Neurology* 2000; **54**: 1510-1517.
115. Neerup JL, Gerstenberg T, Kallestrup EB, Koefoed P, Nordling J, Nielsen JE. Urodynamic evaluation of patients with autosomal dominant pure spastic paraplegia linked to chromosome 2p21-2p24. *J Neurol Neurosurg Psychiatry* 1998; **65**: 693-696.
116. Figlewicz D.A. and Bird T.D.. "Pure" hereditary spastic paraplegia-The story becomes complicated. *Neurology* 1999; **53**: 5-7.
117. Fisher E . and Scambler P. Human haploinsufficiency-one for sorrow, two for joy. *Nature Genet.* 1994; **7**: 5-7.

- 
118. Wilkie AOM . The molecular basis of genetic dominace. *J Med Genet.* 1994; **31**: 89-98.
119. Holtzman DM, Epstein CJ. The molecular genetics of Down syndrome. *Mol Genet Med* 1992; **2**: 105-120.
120. Yamakawa K, Huo Yk, Haendel MA, Huber R, Chen XN, Lyons GE, Korenberg JR. DSCAM: a novel member of the immunoglobulin superfamily maps in a Down syndrome region and is involved in the development of the nervous system. *Hum Mol Genet* 1998; **7**: 227-237.
121. Oliner JD, Kinzler KW, Meltzer PS, George DL, Vogelstein B. Amplification of a gene encoding a p53-associated protein in human sarcomas. *Nature* 1992; **358**: 80-83.
122. Cleary ML. Oncogenic conversion of transcription factors by chromosomal translocations. *Cell* 1991; **66**: 619-622.
123. Rabbitts TH. Translocations, master genes, and differences between the origins of acute and chronic leukemias. *Cell* 1991; **67**:641-644.
124. Trent C, Wood WB, Horvitz HR. A novel dominant transformer allele of the sex-determining gene *her-1* of *Caenorhabditis elegans*. *Genetics* 1988; **120**: 145-157.
125. Miller MW, Duhl DM, Vrieling H, Cordes SP, Ollmann MM, Winkes BM, Barsh GS. Cloning of the mouse *agouti* gene predicts a secreted protein ubiquitously expressed in mice carring the *lethal yellow* mutation. *Genes Dev* 1993; **7**: 1203-1213.
126. Feingold EA, Forget BG. The breakpoint of a large deletion causing hereditary persistence of fetal hemoglobin occurs within an erythroid DNA domain remote from the  $\beta$ -globin gene cluster. *Blood* 1989; **74**: 2178-2186.
127. Rogers S, Wells R, Rechsteiner M. Amino acid sequences common to rapidly degraded proteins: the PEST hypothesis. *Science* 1986; **234**: 364-368.

- 
128. Mango SE, Maine EM, Kimble J. Carboxy-terminal truncation activates *glp-1* protein to specify vulval fates in *Caenorhabditis elegans*. *Nature* 1991; **352**: 811-815.
129. Vogelstein B. A deadly inheritance. *Nature* 1990; **348**: 681-682.
130. Vogelstein B, Kinzler KW. P53 function and dysfunction. *Cell* 1992; **70**: 523-526.
131. Monplaisir N, Merault G, Poyart C, Rhoda MD, Craescu C, Vidaud M, Galacteros F, Blouquit Y, Rosa J. Hemoglobin S Antilles: a variant with lower solubility than hemoglobin S and producing sickle cell disease in heterozygotes. *Proc Natl Acad Sci USA* 1986; **83**: 9363-9367.
132. Owen MC, Brennan SO, Lewis JH, Carrell RW. Mutation of antitrypsin to antithrombin  $\alpha_1$ -antitrypsin Pittsburgh (358 Met→Arg), a fatal bleeding disorder. *N Engl J Med* 1983; **309**: 694-698.
133. Knudson AG. Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 1971; **68**: 820-823.
134. Cavenee WK, Dryja TP, Phillips RA, Benedict WF, Godbout R, Gallie BL, Murphree  
135. AL, Strong LC, White RL. Expression of recessive alleles by chromosomal  
mechanisms  
136. in retinoblastoma. *Nature* 1983; **305**: 779-784.
137. DeChiara TM, Roberston EJ, Efstratiadis A. Parental imprinting of the mouse insulin-like growth factor II gene. *Cell* 1991; **64**: 849-859.